



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Bocconoline

Cat. No.:	HY-150058
CAS No.:	32906-88-0
Molecular Formula:	C ₂₂ H ₂₁ NO ₅
Molecular Weight:	379.41
Target:	Others
Pathway:	Others
Storage:	4°C, protect from light * In solvent : -80°C, 6 months; -20°C, 1 month (protect from light)



SOLVENT & SOLUBILITY

In Vitro

DMSO : 3 mg/mL (7.91 mM; Need ultrasonic and warming)

Concentration	Solvent	Mass		
		1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM	2.6357 mL	13.1784 mL	26.3567 mL
	5 mM	0.5271 mL	2.6357 mL	5.2713 mL
	10 mM	---	---	---

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

Bocconoline is a potent early endosome antigen 1 (EEA1) inhibitor. Bocconoline can be isolated from *Macleaya cordata*. Bocconoline can be used for the research of Parkinson's disease (PD)^[1].

IC₅₀ & Target

EEA1^[1]

In Vitro

Bocconoline (compound 3) (30 μM; 24 h; hONS cells) affects primarily on EEA1 marker intensity in the cytoplasm, mitochondria marker intensity in inner region of the cytoplasm and microtubule-associated protein light chain 3B (LC3b) marker intensity in inner region of cytoplasm^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Nguyen DT, et al. Chemical constituents from *Macleaya cordata* (Willd) R. Br. and their phenotypic functions against a Parkinson's disease patient-derived cell line. *Bioorg Med Chem.* 2020 Nov 1;28(21):115732.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA