



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## HOE-S 785026 trihydrochloride

<b>Cat. No.:</b>	HY-15561B
<b>CAS No.:</b>	2320308-12-9
<b>Molecular Formula:</b>	C <sub>25</sub> H <sub>27</sub> Cl <sub>3</sub> N <sub>6</sub> O
<b>Molecular Weight:</b>	533.88
<b>Target:</b>	Fluorescent Dye
<b>Pathway:</b>	Others
<b>Storage:</b>	4°C, sealed storage, away from moisture and light * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light)



### SOLVENT & SOLUBILITY

#### In Vitro

H<sub>2</sub>O : 47.3 mg/mL (88.60 mM; Need ultrasonic and warming)

Concentration	Solvent	Mass		
		1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM	1.8731 mL	9.3654 mL	18.7308 mL
	5 mM	0.3746 mL	1.8731 mL	3.7462 mL
	10 mM	0.1873 mL	0.9365 mL	1.8731 mL

Please refer to the solubility information to select the appropriate solvent.

### BIOLOGICAL ACTIVITY

#### Description

HOE-S 785026 trihydrochloride is a marker dye in Hoechst series. Hoechst is A live nuclear marker dye. Hoechst binds to the grooves in the DNA double strand, which tends to be A/ T-rich DNA strand. Although it binds to all nucleic acids, the A/ T-rich double strand DNA significantly enhances fluorescence intensity Therefore, Hoechst dye can be used for living cell labeling. The fluorescence intensity of Hoechst dye increases with the increase of pH of solution<sup>[1]</sup>.

#### In Vitro

##### General Protocol

##### Preparation of Hoechst working solution

##### 1.1 Preparation of the stock solution

Dissolve 10 mg of in 5 mL DMSO

Note: It is recommended to store the stock solution at 4° or -20° away from light and avoid repetitive freeze-thaw cycles.

##### 1.2 Preparation of Hoechst working solution

Dilute the stock solution in serum-free cell culture medium or PBS to obtain final concentration 10 µg/mL Hoechst working solution.

Note: Please adjust the concentration of Hoechst working solution according to the actual situation.

##### 1. Cell staining

##### 2.1 Suspension cells 6-well plate

- a. Centrifuge at 1000 g at 4°C for 3-5 minutes and then discard the supernatant. Wash twice with PBS, 5 minutes each time. The cell density is  $1 \times 10^6$ /mL.
- b. Add 1 mL of working solution, and then incubate at room temperature for 3-10 minutes.
- c. Centrifuge at 400 g at 4°C for 3-4 minutes and then discard the supernatant.
- d. Wash twice with PBS, 5 minutes each time.
- e. Resuspend cells with serum-free cell culture medium or PBS. Observation by fluorescence microscopy or flow cytometry.

#### 2.2 Adherent cells

- a. Culture adherent cells on sterile coverslips.
- b. Remove the coverslip from the medium and aspirate excess medium.
- c. Add 100  $\mu$ L of working solution, gently shake it to completely cover the cells, and then incubate at room temperature for 3-10 minutes.
- d. Wash twice with medium, 5 minutes each time. Observation by fluorescence microscopy or flow cytometry.

#### Precautions

1. Please adjust the concentration of Hoechst working solution according to the actual situation.
  2. This product is for R&D use only, not for drug, household, or other uses.
  3. For your safety and health, please wear a lab coat and disposable gloves to operate.
- MCE has not independently confirmed the accuracy of these methods. They are for reference only.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA