



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Kartogenin sodium

Cat. No.:	HY-16268A
CAS No.:	1401168-39-5
Molecular Formula:	C ₂₀ H ₁₄ NNaO ₃
Molecular Weight:	339.32
Target:	TGF-beta/Smad
Pathway:	Stem Cell/Wnt; TGF-beta/Smad
Storage:	4°C, sealed storage, away from moisture and light * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light)



BIOLOGICAL ACTIVITY

Description	Kartogenin (KGN) sodium is an inducer of chondrogenic tissue formation (EC ₅₀ : 100 nM). Kartogenin sodium induces chondrogenesis by binding to fibrin A, disrupting its interaction with the transcription factor core binding factor beta subunit (CBFβ), and by modulating the CBFβ-RUNX1 transcriptional program. Kartogenin sodium also promotes tendon-bone junction (TBJ) wound healing by stimulating collagen synthesis. Kartogenin sodium is widely used in cell-free therapy in the field of regeneration for cartilage regeneration and protection, tendon-bone healing, wound healing and limb development. Kartogenin sodium promotes cartilage repair, coordinates limb development, and is also used in osteoarthritis (OA) research ^{[1][2][3][4]} .
In Vitro	Kartogenin sodium (100 nM; 72 h) induces chondrocyte nodule formation in primary hMSCs ^[1] . Kartogenin sodium (10 nM-10 μM; 72 h) increases chondrocyte-specific gene expression in hMSCs ^[1] . Kartogenin sodium (0.12-10 μM; 48 h) inhibits nitric oxide (NO) and glycosaminoglycan (GAG) release induced by cytokines in primary bovine articular chondrocytes ^[1] . Kartogenin sodium (50-5000 nM; 2 weeks) induces the chondrogenetic differentiation of the BMSCs in a concentration-dependent manner ^[2] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.
In Vivo	Kartogenin sodium (10 μM in 4 μL of saline; i.a. on days 7 and 21) promotes cartilage repair in collagenase VII-induced OA models in mice ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Sci Bull. 2023 Aug 1.
- Chem Eng J. 1 March 2022, 133861.
- Chem Eng J. 400 (2020) 126004.
- Biomaterials. 2022 Jun;285:121530.
- Biomaterials. December 2021, 121216.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Cai J, Zhang L, Chen J, et al. Kartogenin and its application in regenerative medicine[J]. Current medical science, 2019, 39(1): 16-20.
- [2]. Zhang J, Wang J H C. Kartogenin induces cartilage-like tissue formation in tendon–bone junction[J]. Bone research, 2014, 2(1): 1-10.
- [3]. Johnson K, et, al. A stem cell-based approach to cartilage repair. Science. 2012 May 11;336(6082):717-21.
- [4]. Liu F, et, al. A novel kartogenin-platelet-rich plasma gel enhances chondrogenesis of bone marrow mesenchymal stem cells in vitro and promotes wounded meniscus healing in vivo. Stem Cell Res Ther. 2019 Jul 8;10(1):201.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA