



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## IITR08367

Cat. No.:	HY-163473
Molecular Formula:	C <sub>16</sub> H <sub>18</sub> S <sub>2</sub>
Molecular Weight:	274.44
Target:	Bacterial
Pathway:	Anti-infection
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



### BIOLOGICAL ACTIVITY

<b>Description</b>	IITR08367 is a potent effector pump <i>Acinetobacter baumannii</i> Fosfomycin Efflux pump (AbaF) inhibitor for enhancing the antimicrobial activity of Fosfomycin (HY-B1075A) against <i>Acinetobacter baumannii</i> . IITR08367 acts by interfering with Fosfomycin/H <sup>+</sup> reverse transporter activity. <sup>[1]</sup>								
<b>In Vitro</b>	<p>IITR08367 (0-50 μM) concentration-dependently enhances the activity of Fosfomycin against AbaF-expressing strain of <i>Escherichia coli</i>. A concentration of 50 μM can increase the postantibiotic effect of Fosfomycin by 30 min<sup>[1]</sup>.</p> <p>IITR08367 (25 μM; 16 min) disrupts the H<sup>+</sup> gradient across the membrane, ultimately inhibiting the H<sup>+</sup> gradient-driven efflux pump without causing membrane damage<sup>[1]</sup>.</p> <p>IITR08367 (100 μM; 12 h) and Fosfomycin (64 mg/L) in combination can inhibit <i>A. baumannii</i> RPTC-15 growth<sup>[1]</sup>.</p> <p>IITR08367 (100 μM) eliminates the Biofilm Forming Ability of <i>A. baumannii</i> with and without the addition of Fosfomycin<sup>[1]</sup>.</p> <p>IITR08367 (12.5-200 μM) is not toxic to erythrocytes<sup>[1]</sup>.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p> <p>Cell Cytotoxicity Assay<sup>[1]</sup></p> <table border="1"> <tr> <td>Cell Line:</td> <td>RBCs</td> </tr> <tr> <td>Concentration:</td> <td>12.5; 25; 50; 100; 200 μM</td> </tr> <tr> <td>Incubation Time:</td> <td>6 h</td> </tr> <tr> <td>Result:</td> <td>Showed less than 40% cytotoxicity till 200 μM</td> </tr> </table>	Cell Line:	RBCs	Concentration:	12.5; 25; 50; 100; 200 μM	Incubation Time:	6 h	Result:	Showed less than 40% cytotoxicity till 200 μM
Cell Line:	RBCs								
Concentration:	12.5; 25; 50; 100; 200 μM								
Incubation Time:	6 h								
Result:	Showed less than 40% cytotoxicity till 200 μM								
<b>In Vivo</b>	<p>IITR08367 (i.p.; 30 mg/kg; every 12 h for 54 h) co-administered with Fosfomycin to Urinary tract infections mice shows significant improvement in the damage caused by <i>A. baumannii</i> RPTC-15, with tissue ultrastructure similar to that of uninfected mice<sup>[1]</sup>.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p> <table border="1"> <tr> <td>Animal Model:</td> <td>Urinary tract infections mice<sup>[1]</sup></td> </tr> <tr> <td>Dosage:</td> <td>30 mg/kg; every 12 h for 54 h</td> </tr> <tr> <td>Administration:</td> <td>i.p.</td> </tr> </table>	Animal Model:	Urinary tract infections mice <sup>[1]</sup>	Dosage:	30 mg/kg; every 12 h for 54 h	Administration:	i.p.		
Animal Model:	Urinary tract infections mice <sup>[1]</sup>								
Dosage:	30 mg/kg; every 12 h for 54 h								
Administration:	i.p.								

---

Result:	Reduced the load of <i>A. baumannii</i> RPTC-15 in the kidney and bladder by approximately $3\log_{10}$ .
---------	---

---

## REFERENCES

---

[1]. Saini M, et al. Small Molecule IITR08367 Potentiates Antibacterial Efficacy of Fosfomycin against *Acinetobacter baumannii* by Efflux Pump Inhibition. *ACS Infect Dis.* 2024 Apr 1.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA