



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Rivastigmine-d₃ hydrochloride

Cat. No.:	HY-17368S2
Molecular Formula:	C ₁₄ H ₂₀ D ₃ ClN ₂ O ₂
Molecular Weight:	289.82
Target:	Cholinesterase (ChE); Isotope-Labeled Compounds
Pathway:	Neuronal Signaling; Others
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description	Rivastigmine-d ₃ hydrochloride is deuterated labeled Rivastigmine (HY-17368). Rivastigmine (S-Rivastigmine) is an orally active and potent cholinesterase (ChE) inhibitor and inhibits butyrylcholinesterase (BChE) and acetylcholinesterase (AChE) with IC ₅₀ s of 0.037 μM, 4.15 μM, respectively. Rivastigmine can pass the blood brain barrier (BBB). Rivastigmine is a parasympathomimetic or cholinergic agent used for the research of mild to moderate dementia of the Alzheimer's type and dementia due to Parkinson's disease ^{[1][2]} .
In Vitro	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs ^[1] . Rivastigmine (S-Rivastigmine; 1 μM; 24 hours) reduces LPS (2.5 μg/ml)-induced TNF-α and IL-6 by 50% and 46% combined with carbachol (10 μM), respectively and does not cause any significant reduction in pro-inflammatory cytokines ^[4] . Rivastigmine (1 μM), carbachol (10 μM), or a combination of both drugs, does not have a cytotoxic effect on activated cells ^[4] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.
In Vivo	Rivastigmine (S-Rivastigmine; 0.5-2.5 mg/kg; IP; 60 min before the tests) significantly and dose-dependently improved the behavioral impairments caused by Aluminum (HY-B1521) ^[5] . Rivastigmine (0.5, 1 mg/kg/day; s.c; for 8 days) reduces by about 50% and 60% respectively, the concentration of IL-6 but not those of TNF-α and IL-1β in BALB/c OlaHsd male mice aged 8-9 weeks weighing 200-250 g with acute colitis ^[4] . Rivastigmine (1 mg/kg), but not (0.5 mg/kg), partially antagonized colon shrinkage and completely prevented bleeding. Treatment with rivastigmine (0.5 mg/kg) causes little change in these pathological manifestations, but rivastigmine (1 mg/kg) causes a partial restoration of the structure of the crypts and a reduction in sub-mucosal edema and cell infiltration. Rivastigmine (1 mg/kg) causes a 4.7% reduction in body weight at the end of the experiment ^[4] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

- [1]. Qian-Sheng Yu, et al. Anticholinesterase activity of compounds related to geneserine tautomers. N-Oxides and 1,2-oxazines. J Med Chem. 2002 Aug 15;45(17):3684-91.
- [2]. Helena Shifrin, et al. Rivastigmine alleviates experimentally induced colitis in mice and rats by acting at central and peripheral sites to modulate immune responses. PLoS One. 2013;8(2):e57668.

[3]. Han HJ, Lee JJ, Park SA et al. Efficacy and safety of switching from oral cholinesterase inhibitors to the rivastigmine transdermal patch in patients with probable Alzheimer's disease. J Clin Neurol. 2011 Sep;7(3):137-42.

[4]. Raafat A Abdel-Aal, et al. Rivastigmine reverses aluminum-induced behavioral changes in rats. Eur J Pharmacol. 2011 Jun 1;659(2-3):169-76.

[5]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019 Feb;53(2):211-216.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA