



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Edonerpic maleate

|                           |  |
|---------------------------|--|
| <b>Cat. No.:</b>          | HY-17631A  |
| <b>CAS No.:</b>           | 519187-97-4  |
| <b>Molecular Formula:</b> | C <sub>20</sub> H <sub>25</sub> NO <sub>6</sub> S  |
| <b>Molecular Weight:</b>  | 407.48   |
| <b>Target:</b>            | Amyloid- $\beta$   |
| <b>Pathway:</b>           | Neuronal Signaling   |
| <b>Storage:</b>           | 4°C, sealed storage, away from moisture<br>* In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture) |



### SOLVENT & SOLUBILITY

|   |  |                      |             |             |             |              |
|---|--|----------------------|-------------|-------------|-------------|--------------|
| <b>In Vitro</b>   | DMSO : 100 mg/mL (245.41 mM; Need ultrasonic)  |                      |             |             |             |              |
|   | H <sub>2</sub> O : 100 mg/mL (245.41 mM; Need ultrasonic)  |                      |             |             |             |              |
|   | <b>Preparing Stock Solutions</b>   | <b>Solvent</b>       | <b>Mass</b> | <b>1 mg</b> | <b>5 mg</b> | <b>10 mg</b> |
|   |  | <b>Concentration</b> |             |             |             |              |
|   |  | <b>1 mM</b>          |             | 2.4541 mL   | 12.2705 mL  | 24.5411 mL   |
| <b>5 mM</b>   |  |                      | 0.4908 mL   | 2.4541 mL   | 4.9082 mL   |              |
|   | <b>10 mM</b>   |                      | 0.2454 mL   | 1.2271 mL   | 2.4541 mL   |              |
| Please refer to the solubility information to select the appropriate solvent. |  |                      |             |             |             |              |
| <b>In Vivo</b>  | 1. Add each solvent one by one: PBS<br>Solubility: 140 mg/mL (343.58 mM); Clear solution; Need ultrasonic                                  |                      |             |             |             |              |
|   | 2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE- $\beta$ -CD in saline)<br>Solubility: $\geq$ 2.5 mg/mL (6.14 mM); Clear solution |                      |             |             |             |              |
|   | 3. Add each solvent one by one: 10% DMSO >> 90% corn oil<br>Solubility: $\geq$ 2.5 mg/mL (6.14 mM); Clear solution                         |                      |             |             |             |              |

### BIOLOGICAL ACTIVITY

|                                     |  |
|-------------------------------------|--|
| <b>Description</b>                  | Edonerpic maleate is a novel neurotrophic agent which can inhibit amyloid- $\beta$ peptides (A $\beta$ ).  |
| <b>IC<sub>50</sub> &amp; Target</b> | amyloid- $\beta$ peptides <sup>[1]</sup>   |
| <b>In Vitro</b>                     | Edonerpic maleate (T-817MA) treatment preserves the cortical neurons in the presence of A $\beta$ (1-42). Twenty-four hours of pretreatment, followed by the continuous presence of Edonerpic maleate, prevents oxidative stress-induced neuronal death at 0.1 and 1 $\mu$ M. Edonerpic maleate almost completely prevents GSH reduction at 0.1 and 1 $\mu$ M. Hippocampal slices with 1 $\mu$ M Edonerpic maleate treatment generate more and much longer neurites than control slices. Edonerpic maleate |

significantly increases the neurite length at 0.1 and 1  $\mu\text{M}$ <sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### In Vivo

The post hoc test indicates that the mean density of PSA-positive cells is significantly larger in the vehicle and A $\beta$  infusion+high-dose Edonerpic maleate (T-817MA) groups than that in the A $\beta$  infusion control group ( $P < 0.01$ ). The results indicate that the vehicle and A $\beta$  infusion+high-dose Edonerpic maleate groups display efficient learning in the place learning task (PLT), while these two groups also display vigorous neurogenesis. Treatment with Edonerpic maleate and donepezil does not increase the mean density of normal granule cells; there are no significant differences in the mean granule cell density among the A $\beta$  infusion control, A $\beta$  infusion+high-dose Edonerpic maleate, A $\beta$  infusion+low-dose Edonerpic maleate and A $\beta$  infusion+donepezil groups<sup>[2]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## PROTOCOL

#### Cell Assay <sup>[1]</sup>

A cortical neuron/glia coculture is prepared. Edonerpic maleate (T-817MA) is added to the cocultures at concentrations of 0 (control), 0.01, 0.1, and 1  $\mu\text{M}$ , and the cells are subsequently incubated for 5 min or 24 h. H<sub>2</sub>O<sub>2</sub> is then added to the coculture at a concentration of 100  $\mu\text{M}$ , and the cells are incubated for another 24 h. For the normal group, the preparations are maintained in the medium with neither Edonerpic maleate nor H<sub>2</sub>O<sub>2</sub>. Neuronal cell viability is quantified by measuring the Monoclonal anti-microtubule-associated protein 2 (MAP2) immunoreactivity<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### Animal Administration <sup>[2]</sup>

Wistar rats (7 weeks, n=47) are used in this study. All rats are given food and water ad libitum in a clear cage and handled on three consecutive days before start of the experiments. The housing area is provided a temperature-controlled environment under a 12/12 h light cycle. These rats are divided into five groups: vehicle (n=11), A $\beta$  infusion control (n=10), A $\beta$  infusion+high-dose Edonerpic maleate (T-817MA) (8.4 mg/kg) (n=11), A $\beta$  infusion+low-dose Edonerpic maleate (0.84 mg/kg) (n=9) and A $\beta$  infusion+donepezil (0.5 mg/kg) (n=7)<sup>[2]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## REFERENCES

[1]. Hirata K, et al. A novel neurotrophic agent, T-817MA [1-{3-[2-(1-benzothiophen-5-yl) ethoxy] propyl}-3-azetidinol maleate], attenuates amyloid-beta-induced neurotoxicity and promotes neurite outgrowth in rat cultured central nervous system neurons. *J Pharmacol Exp Ther.* 2005 Jul;314(1):252-9.

[2]. Kimura T, et al. T-817MA, a neurotrophic agent, ameliorates the deficits in adult neurogenesis and spatial memory in rats infused i.c.v. with amyloid-beta peptide. *Br J Pharmacol.* 2009 Jun;157(3):451-63.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA