



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## KML29

<b>Cat. No.:</b>	HY-18977		
<b>CAS No.:</b>	1380424-42-9		
<b>Molecular Formula:</b>	C <sub>24</sub> H <sub>21</sub> F <sub>6</sub> NO <sub>7</sub>		
<b>Molecular Weight:</b>	549.42		
<b>Target:</b>	MAGL		
<b>Pathway:</b>	Metabolic Enzyme/Protease		
<b>Storage:</b>	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	2 years
		-20°C	1 year



### SOLVENT & SOLUBILITY

<b>In Vitro</b>	DMSO : 50 mg/mL (91.01 mM; Need ultrasonic)			
		Solvent Concentration	Mass	
			1 mg	5 mg
			10 mg	
<b>Preparing Stock Solutions</b>	<b>1 mM</b>	1.8201 mL	9.1005 mL	18.2010 mL
	<b>5 mM</b>	0.3640 mL	1.8201 mL	3.6402 mL
	<b>10 mM</b>	0.1820 mL	0.9101 mL	1.8201 mL
Please refer to the solubility information to select the appropriate solvent.				
<b>In Vivo</b>	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (4.55 mM); Clear solution  2. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (4.55 mM); Clear solution			

### BIOLOGICAL ACTIVITY

<b>Description</b>	KML29 is an extremely selective, orally active and irreversible MAGL inhibitor, with IC <sub>50</sub> values of 15 nM, 43 nM and 5.9 nM for mouse, rat and human MAGL, respectively. KML29 exhibits minimal cross-reactivity toward other central and peripheral serine hydrolases, including no detectable activity against FAAH <sup>[1][2]</sup> .
<b>IC<sub>50</sub> &amp; Target</b>	IC <sub>50</sub> : 15 nM (mouse MAGL), 43 nM (rat MAGL), 5.9 nM (human MAGL) <sup>[2]</sup> .
<b>In Vitro</b>	KML29 dose-dependently elevates brain 2-AG level up to 10-fold without alteration in brain levels of anandamide, palmitoylethanolamide, and oleoylethanolamide <sup>[2]</sup> . KML29 is a potent inhibitor of 2-AG hydrolysis, but did not affect AEA hydrolysis at any concentration tested <sup>[2]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## In Vivo

KML29 inhibits antinociceptive activity without cannabimimetic side effects<sup>[3]</sup>.  
KML29 (20 mg/kg) has a significant but modest protective effect against LPS-induced fever<sup>[3]</sup>.  
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	C57Bl/6 mice <sup>[2]</sup> .
---------------	-------------------------------

Dosage:	1-40 mg/kg.
---------	-------------

Administration:	P.O. single dose.
-----------------	-------------------

Result:	Selectively inhibited MAGL in mice.
---------	-------------------------------------

Animal Model:	Wistar albino male rats <sup>[2]</sup> .
---------------	--

Dosage:	20 mg/kg (+LPS E. coli O111:B4 (250 µg/kg, sc)).
---------	--

Administration:	SC.
-----------------	-----

Result:	Administration of KML29 simultaneously with LPS E. coli O111:B4 significantly decreased $\Delta T$ (with 5% type 1 error, 1.7 fold) compared to saline+LPS E. coli O111:B4. Administration of KML29 simultaneously with LPS E. coli O111:B4 resulted in decreased plateau phase of fever compared to LPS E. coli O111:B4+saline administration.
---------	---

## CUSTOMER VALIDATION

- Arthritis Res Ther. 2020 Jan 14;22(1):9.
- Dalhousie University. 2017 Nov.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

[1]. Natsuo Ueda, et al. Discrimination between two endocannabinoids. Chem Biol. 2012 May 25;19(5):545-7.

[2]. Jae Won Chang, et al. Highly selective inhibitors of monoacylglycerol lipase bearing a reactive group that is bioisosteric with endocannabinoid substrates. Chem Biol. 2012 May 25;19(5):579-88.

[3]. B M Ignatowska-Jankowska, et al. In vivo characterization of the highly selective monoacylglycerol lipase inhibitor KML29: antinociceptive activity without cannabimimetic side effects. Br J Pharmacol. 2014 Mar;171(6):1392-407.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA