



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

D-Cycloserine

Cat. No.:	HY-B0030												
CAS No.:	68-41-7												
Molecular Formula:	C ₃ H ₆ N ₂ O ₂												
Molecular Weight:	102.09												
Target:	Bacterial; Antibiotic; iGluR												
Pathway:	Anti-infection; Membrane Transporter/Ion Channel; Neuronal Signaling												
Storage:	<table border="0"> <tr> <td>Powder</td> <td>-20°C</td> <td>3 years</td> </tr> <tr> <td></td> <td>4°C</td> <td>2 years</td> </tr> <tr> <td>In solvent</td> <td>-80°C</td> <td>2 years</td> </tr> <tr> <td></td> <td>-20°C</td> <td>1 year</td> </tr> </table>	Powder	-20°C	3 years		4°C	2 years	In solvent	-80°C	2 years		-20°C	1 year
Powder	-20°C	3 years											
	4°C	2 years											
In solvent	-80°C	2 years											
	-20°C	1 year											



SOLVENT & SOLUBILITY

In Vitro	H ₂ O : 25 mg/mL (244.88 mM; Need ultrasonic)				
	DMSO : 1 mg/mL (9.80 mM; Need warming)				
		Solvent Concentration	Mass		
	Preparing Stock Solutions		1 mg	5 mg	10 mg
	1 mM	9.7953 mL	48.9764 mL	97.9528 mL	
	5 mM	1.9591 mL	9.7953 mL	19.5906 mL	
	10 mM	0.9795 mL	4.8976 mL	9.7953 mL	
Please refer to the solubility information to select the appropriate solvent.					
In Vivo	1. Add each solvent one by one: PBS Solubility: 11.11 mg/mL (108.83 mM); Clear solution; Need ultrasonic				

BIOLOGICAL ACTIVITY

Description	D-Cycloserine is an antibiotic which targets sequential bacterial cell wall peptidoglycan biosynthesis enzymes. D-Cycloserine is a partial NMDA agonist that can improve cognitive functions. D-Cycloserine can be used for multidrug-resistant tuberculosis research ^{[1][2]} .
IC₅₀ & Target	NMDA Receptor
In Vivo	D-Cycloserine selectively potentiates the duration of motor cortical excitability enhancements induced by anodal transcranial direct current (tDCS). D-Cycloserine alone does not modulate excitability. ^[1] Chronic D-Cycloserine (40 mg/kg; 5 days/week; for 2 weeks) significantly reduces nicotine self-administration selectively in rats with low baseline nicotine use, but is ineffective with the rats with higher levels of baseline nicotine self-administration ^[3] .

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- ACS Chem Biol. 2021 Dec 15.
- Exp Brain Res. 2023 Oct 16.
- Research Square Preprint. 2021 Aug.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Nitsche, M.A., et al., Consolidation of human motor cortical neuroplasticity by D-cycloserine. *Neuropsychopharmacology*, 2004. 29(8): p. 1573-8.
- [2]. Levin, E.D., et al., D-cycloserine selectively decreases nicotine self-administration in rats with low baseline levels of response. *Pharmacol Biochem Behav*, 2011. 98(2): p. 210-4.
- [3]. Sarah Batson, et al. Inhibition of D-Ala:D-Ala ligase through a phosphorylated form of the antibiotic D-cycloserine. *Nat Commun*. 2017 Dec 5;8(1):1939.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA