



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Moxonidine-¹³C,₃D₃ hydrochloride

Cat. No.:	HY-B0374AS
Molecular Formula:	C ₈ ¹³ CH ₁₀ D ₃ Cl ₂ N ₅ O
Molecular Weight:	282.15
Target:	Imidazoline Receptor; Isotope-Labeled Compounds
Pathway:	Neuronal Signaling; Others
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description	Moxonidine- ¹³ C, ₃ D ₃ hydrochloride is ¹³ C and deuterated labeled Moxonidine hydrochloride (HY-B0374A). Moxonidine (BDF5895) hydrochloride is an imidazoline type 1 receptor (I1-R) selective agonist and antihypertensive agent.
In Vitro	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

- [1]. Schafer, U., et al., Presynaptic effects of moxonidine in isolated buffer perfused rat hearts: role of imidazoline-1 receptors and alpha2-adrenoceptors. *J Pharmacol Exp Ther*, 2002. 303(3): p. 1163-70.
- [2]. Chan, C.K., et al., Imidazoline receptors associated with noradrenergic terminals in the rostral ventrolateral medulla mediate the hypotensive responses of moxonidine but not clonidine. *Neuroscience*, 2005. 132(4): p. 991-1007.
- [3]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. *Ann Pharmacother*. 2019 Feb;53(2):211-216.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA