

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



Hydralazine-d₅ hydrochloride


Cat. No.: HY-B0464S1 CAS No.: 2012598-61-5 Molecular Formula: C₈H₄D₅ClN₄

Molecular Weight: 201.67

Target: Caspase; Bcl-2 Family; Apoptosis; Isotope-Labeled Compounds

Pathway: Apoptosis; Others

Storage: Please store the product under the recommended conditions in the Certificate of



Product Data Sheet

BIOLOGICAL ACTIVITY

Description

Hydralazine-d₅ (hydrochloride) is deuterium-labeled Hydralazine (hydrochloride) (HY-B0464)^[1].

In Vitro

Stable or radioisotope-labeled compounds allow precise tracking and quantification of individual atoms in metabolic pathways. Stable isotopes generally do not change molecular properties but may slightly affect metabolic kinetics; radioactive isotopes may interfere with cells. Markers can distinguish endogenous and exogenous metabolites, reduce false positives, and are beneficial to quantification and reconstruction of metabolic pathways^[2].

In cell culture or enzymatic reactions, the use of isotope markers can precisely control the concentration and exposure time, making it easy to study metabolic reactions and enzyme activities. Through stable isotope analytical metabolomics (SIRM), cellular metabolic networks can be studied, key metabolic nodes and regulatory mechanisms can be identified, and targets can be provided for compound development.

Isotope-labeled compounds can be used in competition binding experiments to evaluate the affinity and binding kinetics of compounds to receptors to help optimize design. Stable isotope labels are used as internal standards in mass spectrometry analysis to improve analysis accuracy and reproducibility and reduce matrix effect interference^[3].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

Isotopic labels can non-invasively track the distribution, transformation and clearance of compounds and their metabolites in the body through techniques such as mass spectrometry (MS) and nuclear magnetic resonance (NMR), which is beneficial to the study of pharmacometabolic kinetics (ADME).

Isotope labeling can reveal specific steps in metabolic pathways. Using compounds with stable isotope labels at specific locations directly in humans or animal models can also help verify drug mechanisms and evaluate unexpected side effects, improving the accuracy and efficiency of clinical research^[3].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

- [1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019;53(2):211-216.
- [2]. Smith K A, et al. Soil and environmental analysis[M]. Marcel Dekker Incorporated, 2000.
- [3]. Fan T W M, et al. Stable isotope-resolved metabolomics and applications for drug development[J]. Pharmacology & therapeutics, 2012, 133(3): 366-391.

 $\label{lem:caution:Product} \textbf{Caution: Product has not been fully validated for medical applications. For research use only.}$

Tel: 609-228-6898 Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

Page 2 of 2 www.MedChemExpress.com