



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Nifursol-<sup>13</sup>C<sub>6</sub>

<b>Cat. No.:</b>	HY-B1703S
<b>Molecular Formula:</b>	C <sub>6</sub> <sup>13</sup> C <sub>6</sub> H <sub>7</sub> N <sub>5</sub> O <sub>9</sub>
<b>Molecular Weight:</b>	371.17
<b>Target:</b>	Bacterial; Antibiotic
<b>Pathway:</b>	Anti-infection
<b>Storage:</b>	4°C, sealed storage, away from moisture and light * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light)



### BIOLOGICAL ACTIVITY

<b>Description</b>	Nifursol- <sup>13</sup> C <sub>6</sub> is the <sup>13</sup> C <sub>6</sub> labeled Nifursol. Nifursol is a potent and orally active veterinary antibiotic for the prevention of histomoniasis. Nifursol rapidly metabolizes to form the metabolic marker 3,5-dinitrosalicylic acid hydrazide (DNSAH) which can persist for a long time. Nifursol is widely used for the research of Escherichia Gastroenteropathy in poultry, fowl and aquatic animal.
<b>In Vitro</b>	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### REFERENCES

- [1]. Bowen TE, et al. Effect of cupric sulfate on the prophylactic efficacy of 2-acetylamino-5nitrothiazole, nifursol and ipronidazole against histomoniasis in turkeys. *Poult Sci.* 1971 Nov;50(6):1668-72.
- [2]. Jorge Barbosa, et al. Detection, accumulation, distribution, and depletion of furaltadone and nifursol residues in poultry muscle, liver, and gizzard. *J Agric Food Chem.* 2011 Nov 23;59(22):11927-34.
- [3]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. *Ann Pharmacother.* 2019 Feb;53(2):211-216.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA