

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



Proteins

Product Data Sheet

Chondroitin sulfate sodium

Cat. No.: HY-B2162A CAS No.: 9082-07-9

Target: NO Synthase; MMP; Endogenous Metabolite

Pathway: Immunology/Inflammation; Metabolic Enzyme/Protease

4°C, sealed storage, away from moisture and light Storage:

* In solvent: -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture

and light)

 $R_2 = H$ R₂= SO₃H $R_1 = SO_3H$

SOLVENT & SOLUBILITY

In Vitro

H₂O: 100 mg/mL (Need ultrasonic)

BIOLOGICAL ACTIVITY

_			
Desc	rıp	tioi	n

Chondroitin sulfate sodium, one of five classes of glycosaminoglycans, has been widely used in the treatment of osteoarthritis. Chondroitin sulfate reduces inflammation mediators and the apoptotic process and is able to reduce protein production of inflammatory cytokines, iNOS and MMPs^{[1][2][3][4]}.

IC₅₀ & Target

iNOS

Human Endogenous Metabolite

In Vitro

Chondroitin sulfate is a class of sulfated glycosaminoglycans that are linear polysaccharides consisting of repeating disaccharide units composed of uronic acid and N-acetylhexosamine. Several pathogens including parasites, bacteria, and viruses have been shown to utilize cell surface chondroitin sulfate chains to attach to and infect host cells^[1]. Chondroitin sulfate occurs naturally in the extracellular matrix of connective tissues, e.g., bone, cartilage, skin, ligaments and tendons. Chondroitin sulfate has been shown to elicit a range of beneficial effects: anti-inflammatory effects, an increase in type II collagen and proteoglycans, a reduction in bone resorption and a better anabolic/catabolic balance in chondrocytes^[2].

A large range of chondroitin sulfate concentrations has been used (e.g. 12.5 to 2000 mg/mL, but generally less than 200 mg/mL) in in vitro studies. Chondroitin sulfate (200 mg/mL) decreases the chondrocyte susceptibility to single nucleotide polymorphism-induced apoptosis^[3].

Chondroitin sulfate reduces inflammation mediators and the apoptotic process and is able to reduce protein production of inflammatory cytokines, iNOS, MMPs^[4].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

The high content of chondroitin sulfate in the aggrecan plays a major part in allowing cartilage to resist tensile stresses during various loading conditions by providing this tissue with resistance and elasticity. It has been shown that chondroitin sulphate interferes with the progression of structural changes in joint tissues and is used in the management of patients with osteoarthritis[3].

Chondroitin sulfate is mostly administered orally at doses ranging from 800 to 1200mg/day. Chondroitin sulfate is rapidly absorbed by the gastrointestinal tract. The absorbed chondroitin sulfate reaches the blood compartment as 10% chondroitin sulfate and 90% depolymerized low-molecular-weight derivatives^[5].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Immunity. 2021 May 11;54(5):962-975.e8.
- · Acta Pharm Sin B. 2023 Dec 16.
- J Thromb Haemost. 2021 Feb;19(2):470-477.
- FEBS Lett. 2020 Jan;594(1):94-103.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Mikami T, et al. Biosynthesis and function of chondroitin sulfate. Biochim Biophys Acta. 2013 Oct;1830(10):4719-33.
- [2]. Monfort J, et al. Biochemical basis of the effect of chondroitin sulphate on osteoarthritis articular tissues. Ann Rheum Dis. 2008 Jun;67(6):735-40.
- [3]. Martel-Pelletier J, et al.Discrepancies in composition and biological effects of different formulations of chondroitin sulfate. Molecules. 2015 Mar 6;20(3):4277-89.
- [4]. Campo GM, et al. Glycosaminoglycans modulate inflammation and apoptosis in LPS-treated chondrocytes. J Cell Biochem. 2009 Jan 1;106(1):83-92.
- [5]. Henrotin Y, et al. Chondroitin sulfate in the treatment of osteoarthritis: from in vitro studies to clinical recommendations. Ther Adv Musculoskelet Dis. 2010 Dec;2(6):335-48

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898 Fax: 609-228-5909 E-mail: tech@MedChemExpress.com Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA