



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Quinine

Cat. No.:	HY-D0143
CAS No.:	130-95-0
Molecular Formula:	C ₂₀ H ₂₄ N ₂ O ₂
Molecular Weight:	324.42
Target:	Potassium Channel; Parasite; Flavivirus; Dengue virus
Pathway:	Membrane Transporter/Ion Channel; Anti-infection
Storage:	4°C, protect from light, stored under nitrogen * In solvent : -80°C, 6 months; -20°C, 1 month (protect from light, stored under nitrogen)



SOLVENT & SOLUBILITY

In Vitro

DMSO : ≥ 100 mg/mL (308.24 mM)
 H₂O : < 0.1 mg/mL (insoluble)
 * "≥" means soluble, but saturation unknown.

	Solvent Concentration	Mass		
		1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM	3.0824 mL	15.4121 mL	30.8242 mL
	5 mM	0.6165 mL	3.0824 mL	6.1648 mL
	10 mM	0.3082 mL	1.5412 mL	3.0824 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: ≥ 2.5 mg/mL (7.71 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
Solubility: ≥ 2.5 mg/mL (7.71 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil
Solubility: ≥ 2.5 mg/mL (7.71 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

Quinine is an alkaloid derived from the bark of the cinchona tree, acts as an anti-malaria agent. Quinine is a potassium channel inhibitor that inhibits WT mouse Slo3 (K_{Ca}5.1) channel currents evoked by voltage pulses to +100 mV with an IC₅₀ of 169 μM^{[1][2]}.

IC₅₀ & Target

Plasmodium

In Vitro	<p>Quinine (150 μM, 30 min) inhibits the proliferation and cytostatic effects of DENV (Dengue virus) in human hepatocarcinoma HepG2 cell line^[1].</p> <p>Quinine (37.5-150 μM, 24 hours) significantly reduces viral DENV RNA and protein levels in a dose-dependent manner in human hepatocarcinoma HepG2 cell line^[1].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p> <p>Cell Proliferation Assay^[1]</p>								
	<table border="1"> <tr> <td>Cell Line:</td> <td>Human hepatocarcinoma cell line (HepG2)</td> </tr> <tr> <td>Concentration:</td> <td>150 μM</td> </tr> <tr> <td>Incubation Time:</td> <td>30 min</td> </tr> <tr> <td>Result:</td> <td>Inhibited DENV virus replication with 19% yield compared to untreated. Reduced DENV-positive cells from 23.28% to 12.05% in a dose-dependent manner.</td> </tr> </table>	Cell Line:	Human hepatocarcinoma cell line (HepG2)	Concentration:	150 μ M	Incubation Time:	30 min	Result:	Inhibited DENV virus replication with 19% yield compared to untreated. Reduced DENV-positive cells from 23.28% to 12.05% in a dose-dependent manner.
	Cell Line:	Human hepatocarcinoma cell line (HepG2)							
	Concentration:	150 μ M							
	Incubation Time:	30 min							
Result:	Inhibited DENV virus replication with 19% yield compared to untreated. Reduced DENV-positive cells from 23.28% to 12.05% in a dose-dependent manner.								
In Vivo	<p>Quinine (oral gavage, 12 or 15 mg/kg, every week, 16 weeks) has some tumor suppressing effect on skin cancer in Swiss albino mice^[2].</p> <p>Quinine (oral gavage, 10 mg/kg, everyday, 8 weeks) causes a decrease in the antioxidant defense system of rat testicular tissue such as SOD, CAT and GSH enzyme activity in male adult albino rats^[3].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>								
	<table border="1"> <tr> <td>Animal Model:</td> <td>Swiss albino mice 7-8-weeks (weighing 24 g)^[2]</td> </tr> <tr> <td>Dosage:</td> <td>12 mg/kg, 15 mg/kg</td> </tr> <tr> <td>Administration:</td> <td>Oral gavage; every week; 16 weeks</td> </tr> <tr> <td>Result:</td> <td>Resulted in a significant reduction in tumor size and weight at 12 mg/kg and little effect at higher dose of 15 mg/kg.</td> </tr> </table>	Animal Model:	Swiss albino mice 7-8-weeks (weighing 24 g) ^[2]	Dosage:	12 mg/kg, 15 mg/kg	Administration:	Oral gavage; every week; 16 weeks	Result:	Resulted in a significant reduction in tumor size and weight at 12 mg/kg and little effect at higher dose of 15 mg/kg.
	Animal Model:	Swiss albino mice 7-8-weeks (weighing 24 g) ^[2]							
	Dosage:	12 mg/kg, 15 mg/kg							
	Administration:	Oral gavage; every week; 16 weeks							
Result:	Resulted in a significant reduction in tumor size and weight at 12 mg/kg and little effect at higher dose of 15 mg/kg.								

CUSTOMER VALIDATION

- ACS Omega. 2024 Feb 28;9(10):11870-11882.
- Mol Med Rep. 2021 Mar 2.
- Norwegian University of Science and Technology, Faculty of Medicine and Health sciences. 2019 Sep.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Shilu Malakar et al. Drug repurposing of quinine as antiviral against dengue virus infection. *Virus Res.* 2018 Aug 15;255:171-178. doi: 10.1016/j.virusres.2018.07.018. Epub 2018 Jul 25.
- [2]. Jhanwar, Deepika et al. Chemoprevention of DMBA induced skin carcinogenesis in swiss albino mice by quinine sulfate.(2016): 2636-2640.
- [3]. Ebenezer O Farombi, et al. Quercetin protects against testicular toxicity induced by chronic administration of therapeutic dose of quinine sulfate in rats. *J Basic Clin Physiol Pharmacol.* 2012 Feb 27;23(1):39-44.
- [4]. Wrighton DC, et al. Mechanism of inhibition of mouse Slo3 (KCa 5.1) potassium channels by quinine, quinidine and barium. *Br J Pharmacol.* 2015 Sep;172(17):4355-63.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA