



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Naphthoresorcinol

Cat. No.:	HY-D0165
CAS No.:	132-86-5
Molecular Formula:	C ₁₀ H ₈ O ₂
Molecular Weight:	160.17
Target:	Fluorescent Dye
Pathway:	Others
Storage:	4°C, protect from light * In solvent : -80°C, 6 months; -20°C, 1 month (protect from light)



SOLVENT & SOLUBILITY

In Vitro	DMSO : 100 mg/mL (624.34 mM; Need ultrasonic)				
		Solvent Concentration	Mass		
	Preparing Stock Solutions		1 mg	5 mg	10 mg
		1 mM	6.2434 mL	31.2168 mL	62.4337 mL
		5 mM	1.2487 mL	6.2434 mL	12.4867 mL
	10 mM	0.6243 mL	3.1217 mL	6.2434 mL	
Please refer to the solubility information to select the appropriate solvent.					
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (15.61 mM); Clear solution				
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (15.61 mM); Clear solution				

BIOLOGICAL ACTIVITY

Description	Naphthoresorcinol (1,3-Dihydroxynaphthalene) is a fluorescent dye (λ_{ex} =330 nm, λ_{em} =380 nm) that can react with the NPPD (a tracer) and concentrated HCl and develop a red color. Naphthoresorcinol could be used as a background electrolyte (BGE) to determine the carbohydrates ^{[1][2][3]} .
-------------	---

REFERENCES

- [1]. Vyas S, et, al. Fluorescence and light scattering studies on the interaction of 1,3-dihydroxynaphthalene with ionic and non-ionic surfactants. *Polymer Photochemistry*. 1984; 4(4):245-253.
- [2]. Suzuki S, et, al. Development of a field kit for use by non-scientists for chemical tracking using 5-(4-nitrophenyl)-2,4-pentadien-1-al. *Forensic Sci Int*. 2013 May 10;228(1-3):e25-7.

[3]. Lee YH, et, al. Determination of carbohydrates by high-performance capillary electrophoresis with indirect absorbance detection. J Chromatogr B Biomed Appl. 1996 May 31;681(1):87-97.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA