



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Fluo-3AM

<b>Cat. No.:</b>	HY-D0716
<b>CAS No.:</b>	121714-22-5
<b>Molecular Formula:</b>	C <sub>51</sub> H <sub>50</sub> Cl <sub>2</sub> N <sub>2</sub> O <sub>23</sub>
<b>Molecular Weight:</b>	1129.85
<b>Target:</b>	Fluorescent Dye
<b>Pathway:</b>	Others
<b>Storage:</b>	-20°C, protect from light * In solvent : -80°C, 6 months; -20°C, 1 month (protect from light)



### SOLVENT & SOLUBILITY

#### In Vitro

DMSO : 12.5 mg/mL (11.06 mM; Need ultrasonic)

Concentration	Solvent	Mass		
		1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM	0.8851 mL	4.4254 mL	8.8507 mL
	5 mM	0.1770 mL	0.8851 mL	1.7701 mL
	10 mM	0.0885 mL	0.4425 mL	0.8851 mL

Please refer to the solubility information to select the appropriate solvent.

### BIOLOGICAL ACTIVITY

#### Description

Fluo-3 AM is a fluorescent Ca<sup>2+</sup> chelator, with high affinity for calcium. Fluo-3 AM can specifically identify intracellular calcium ions, with high sensitivity, low cytotoxicity, increased AM acetylmethyl ester can enter the cell well, after being sheared by the intracellular esterase stay in the cell to bind to calcium ions, produce strong fluorescence<sup>[1]</sup>.

#### In Vitro

1. Preparation of Fluo-3 AM working solution
  - 1.1 Preparation of the stock solution  
Dissolve 1 mg Fluo-3 AM in 135 µL DMSO to obtain 10 mM of stock solution.  
Note: It is recommended to store the stock solution at -20°C or -80°C away from light and avoid repetitive freeze-thaw cycles.
  - 1.2 Preparation of Fluo-3 AM working solution  
Dilute the stock solution in HBSS to obtain 1-10 µM of working solution.  
Note: Please adjust the concentration of Fluo-3 AM working solution according to the actual situation.
2. Cell staining
  - 2.1 Suspension cells (6-well plate)
    - a. Centrifuge at 1000 g at 4°C for 3-5 minutes and then discard the supernatant. Wash twice with PBS, 5 minutes each time. The cell density is 1×10<sup>6</sup>/mL
    - b. Add 1 mL of working solution, and then incubate at room temperature for 5-30 minutes.
    - c. Centrifuge at 400 g at 4°C for 3-4 minutes and then discard the supernatant.

- d. Wash twice with PBS, 5 minutes each time.
- e. Resuspend cells with serum-free cell culture medium or PBS. Observation by fluorescence microscopy or flow cytometry.
- 2.2 Adherent cells
- a. Culture adherent cells on sterile coverslips.
- b. Remove the coverslip from the medium and aspirate excess medium.
- c. Add 100  $\mu$ L of working solution, gently shake it to completely cover the cells, and then incubate at room temperature for 5-30 minutes.
- d. Wash twice with medium, 5 minutes each time. Observation by fluorescence microscopy.

Storage

-20°C, 1 year

Protect from light

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## CUSTOMER VALIDATION

- Environ Sci Technol. 2017 Dec 5;51(23):13938-13948.
- Int J Biol Macromol. 2022 Jul 27;S0141-8130(22)01646-4.
- Biomedicines. 2021 Apr 21;9(5):444.
- RSC Adv. 2019, 2019, 9, 25107-25118.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

[1]. Loughrey, C. M., MacEachern, K. E., Cooper, J., & Smith, G. L. (2003). Measurement of the dissociation constant of Fluo-3 for Ca<sup>2+</sup> in isolated rabbit cardiomyocytes using Ca<sup>2+</sup> wave characteristics. Cell Calcium, 34(1), 1–9. doi:10.1016/s0143-4160(03)00012-5

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA