



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Lucigenin

<b>Cat. No.:</b>	HY-D0720
<b>CAS No.:</b>	2315-97-1
<b>Molecular Formula:</b>	C <sub>28</sub> H <sub>22</sub> N <sub>4</sub> O <sub>6</sub>
<b>Molecular Weight:</b>	510.5
<b>Target:</b>	Reactive Oxygen Species; Fluorescent Dye
<b>Pathway:</b>	Immunology/Inflammation; Metabolic Enzyme/Protease; NF-κB; Others
<b>Storage:</b>	4°C, sealed storage, away from moisture and light * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light)



### SOLVENT & SOLUBILITY

<b>In Vitro</b>	DMSO : 62.5 mg/mL (122.43 mM; Need ultrasonic)				
		Solvent Concentration	Mass 1 mg	5 mg	10 mg
	<b>Preparing Stock Solutions</b>	1 mM	1.9589 mL	9.7943 mL	19.5886 mL
		5 mM	0.3918 mL	1.9589 mL	3.9177 mL
		10 mM	0.1959 mL	0.9794 mL	1.9589 mL
Please refer to the solubility information to select the appropriate solvent.					
<b>In Vivo</b>	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.08 mg/mL (4.07 mM); Clear solution  2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.08 mg/mL (4.07 mM); Clear solution				

### BIOLOGICAL ACTIVITY

<b>Description</b>	Lucigenin is a chemiluminescence probe. Lucigenin can be used to detect the production of endogenous superoxide anion radical (O <sup>2-</sup> ). Lucigenin is extremely sensitive to chloride ions, while it combined with chloride ions, the fluorescence will be quenched. Lucigenin also can be used as a chloride indicator. Ex/Em=455/505 nm <sup>[1]</sup> .
<b>In Vitro</b>	Preparation of Lucigenin working solution 1.1 Preparation of the stock solution Dissolve 1 mg of Lucigenin in 0.1919 mL of DMSO to obtain 10 mM of Lucigenin. Note: It is recommended to store the stock solution at -20 °C -80 °C away from light and avoid repetitive freeze-thaw cycles. 1.2 Preparation of Lucigenin working solution Dilute the stock solution in serum-free cell culture medium or PBS to obtain 5-10 μM of Lucigenin working solution. Note: Please adjust the concentration of Lucigenin working solution according to the actual situation.

## Cell staining

### 2.1 Cell preparation.

For suspension cells: Centrifuge at 1000 g at 4°C for 3-5 minutes and then discard the supernatant. Wash twice with PBS, 5 minutes each time.

For adherent cells: Discard the cell culture medium, and add trypsin to dissociate cells to make a single-cell suspension. Centrifuge at 1000 g at 4°C for 3-5 minutes and then discard the supernatant. Wash twice with PBS, 5 minutes each time.

2.2 Add 1 mL of Lucigenin working solution, and then incubate at room temperature for 15 minutes.

2.3 Centrifuge at 400 g at 4°C for 3-4 minutes and then discard the supernatant.

2.4 Wash twice with PBS, 5 minutes each time.

2.5 Resuspend cells with serum-free cell culture medium or PBS, and then detect by fluorescence microscope.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## CUSTOMER VALIDATION

- J Transl Med. 2023 Mar 25;21(1):218.
- Int J Mol Med. 2017 Dec;40(6):1803-1817.
- J Fungi (Basel). 2021 Nov 11;7(11):955.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

[1]. Li Y, et al. Validation of lucigenin (bis-N-methylacridinium) as a chemilumigenic probe for detecting superoxide anion radical production by enzymatic and cellular systems. J Biol Chem. 1998 Jan 23;273(4):2015-23.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA