



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## N-Ethylmaleimide-d<sub>5</sub>

<b>Cat. No.:</b>	HY-D0843S		
<b>CAS No.:</b>	360768-37-2		
<b>Molecular Formula:</b>	C <sub>6</sub> H <sub>2</sub> D <sub>5</sub> NO <sub>2</sub>		
<b>Molecular Weight:</b>	130.16		
<b>Target:</b>	Deubiquitinase; Cathepsin		
<b>Pathway:</b>	Cell Cycle/DNA Damage; Metabolic Enzyme/Protease		
<b>Storage:</b>	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



### SOLVENT & SOLUBILITY

#### In Vitro

DMSO : 100 mg/mL (768.29 mM; Need ultrasonic)

Preparing Stock Solutions	Solvent Concentration	Mass		
		1 mg	5 mg	10 mg
	1 mM	7.6829 mL	38.4143 mL	76.8285 mL
	5 mM	1.5366 mL	7.6829 mL	15.3657 mL
	10 mM	0.7683 mL	3.8414 mL	7.6829 mL

Please refer to the solubility information to select the appropriate solvent.

### BIOLOGICAL ACTIVITY

#### Description

N-Ethylmaleimide-d<sub>5</sub> is the deuterium labeled N-Ethylmaleimide. N-Ethylmaleimide (NEM), a reagent that alkylates free sulfhydryl groups, is a cysteine protease inhibitor[1]. N-ethylmaleimide specific inhibits phosphate transport in mitochondria[2]. N-Ethylmaleimide is also a deubiquitinating enzyme inhibitor[3].

#### In Vitro

Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### REFERENCES

[1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. *Ann Pharmacother.* 2019;53(2):211-216.

[2]. Wu KH, et al. Cys32 and His105 are the critical residues for the calcium-dependent cysteine proteolytic activity of CvaB, an ATP-binding cassette transporter. *J Biol Chem.* 2004;279(12):11511-11517.

---

Chem. 2004 Jan 9;279(2):901-9.

[3]. Hatase O , et al. Specific inhibition of phosphate transport in mitochondria by N-ethylmaleimide. Journal of Bioenergetics, 1973, 5(1):1-15.

[4]. Choo YS, et al. Detection of protein ubiquitination. J Vis Exp. 2009;(30):1293. Published 2009 Aug 19. doi:10.3791/1293

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA