



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

DiR

Cat. No.:	HY-D1048
CAS No.:	100068-60-8
Molecular Formula:	C ₆₃ H ₁₀₁ N ₂
Molecular Weight:	1013.39
Target:	Fluorescent Dye
Pathway:	Others
Storage:	4°C, sealed storage, away from moisture and light * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light)



SOLVENT & SOLUBILITY

In Vitro	DMSO : 25 mg/mL (24.67 mM); ultrasonic and warming and heat to 60°C																					
	<table border="1"> <thead> <tr> <th rowspan="2">Solvent</th> <th rowspan="2">Mass</th> <th colspan="3">Concentration</th> </tr> <tr> <th>1 mg</th> <th>5 mg</th> <th>10 mg</th> </tr> </thead> <tbody> <tr> <td rowspan="3">Preparing Stock Solutions</td> <td>1 mM</td> <td>0.9868 mL</td> <td>4.9339 mL</td> <td>9.8679 mL</td> </tr> <tr> <td>5 mM</td> <td>0.1974 mL</td> <td>0.9868 mL</td> <td>1.9736 mL</td> </tr> <tr> <td>10 mM</td> <td>0.0987 mL</td> <td>0.4934 mL</td> <td>0.9868 mL</td> </tr> </tbody> </table>	Solvent	Mass	Concentration			1 mg	5 mg	10 mg	Preparing Stock Solutions	1 mM	0.9868 mL	4.9339 mL	9.8679 mL	5 mM	0.1974 mL	0.9868 mL	1.9736 mL	10 mM	0.0987 mL	0.4934 mL	0.9868 mL
Solvent	Mass			Concentration																		
		1 mg	5 mg	10 mg																		
Preparing Stock Solutions	1 mM	0.9868 mL	4.9339 mL	9.8679 mL																		
	5 mM	0.1974 mL	0.9868 mL	1.9736 mL																		
	10 mM	0.0987 mL	0.4934 mL	0.9868 mL																		
	Please refer to the solubility information to select the appropriate solvent.																					
In Vivo	1. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.08 mg/mL (2.05 mM); Clear solution																					

BIOLOGICAL ACTIVITY

Description	DiR is a long-chain carbocyanine dye. Carbocyanine dyes are widely used as Di to label cells, organelles, liposomes, viruses and lipoproteins ^[2] .
In Vitro	Carbocyanine dyes are widely used as Di to label cells, organelles, liposomes, viruses and lipoproteins. Long-chain carbocyanines which include DiO (DiOC18(3)), DiI (DiIC18(3)), DiD (DiIC18(5)) and DiR, and dialkyl aminostyryl dye DiA (4-Di-16-ASP) are used for labeling membranes and other hydrophobic structures. DiIC16(3) has shorter alkyl substituents (C16) than DiI (C18). They have extremely high extinction coefficients, environmental dependent fluorescence and short excited-state lifetimes in lipid environments. They are oils at room temperature and weakly fluorescent in water but highly fluorescent and quite photostable when incorporated into membranes or bound to lipophilic biomolecules. These optical characteristics make them ideal for staining the cytoplasmic membranes of cells. Once applied to cells, these dyes diffuse laterally within the plasma membrane, resulting in staining of the entire cell ^[1] . DiO, DiI, DiD and DiR exhibit distinct green, orange, red and infrared fluorescence, respectively thus facilitating multicolor imaging and flow cytometric analysis of live cells . DiO and DiI can be used with standard FITC and TRITC filters respectively.

Among them DiI and its analogs are most frequently used since they usually exhibit very low cell toxicity. In addition, DiI is widely used for determining lipoproteins such as LDL and HDL. The lipophilic aminostyryl dye DiA is also often used for neuronal tracing^[2].

General Protocol

1. Preparing Stain Solutions of Di

a. Prepare DMF, DMSO or ethanol stock solutions: The stock solutions should be prepared in dimethyl formamide (DMF), dimethylsulfoxide (DMSO, or ethanol DMSO at 1-5 mM. DMF is preferable to ethanol as a solvent for Di. The stock solution should be used promptly. Any unused solution need to be aliquoted and refrozen at least -20°C. Avoid repeated freeze/thaw cycle. The solution can be stored for 6 months.

b. Prepare working solutions: Dilute the stock solutions into a suitable buffer such as serum-free culture medium, HBSS or PBS to make 1 to 5 µM working solutions. We do not recommend storing the aqueous solution for more than one day.

Note: The final concentration of the working solution should be empirically determined for different cell types and/or experimental conditions.

2. Suspension cells

a. Centrifuge at 1000 g at 4°C for 3-5 minutes and then discard the supernatant. Wash twice with PBS, 5 minutes each time. The cell density is 1×10^6 /mL.

b. Add 1 mL of Di working solution, and then incubate at room temperature for 5-30 minutes.

c. Centrifuge at 400 g at 4°C for 3-4 minutes and then discard the supernatant.

d. Wash twice with PBS, 5 minutes each time.

e. Resuspend cells with serum-free cell culture medium or PBS. Observation by fluorescence microscopy or flow cytometry.

3. Adherent cells

a. Culture adherent cells on sterile coverslips.

b. Remove the coverslip from the medium and aspirate excess medium.

c. Add 100 µL of working solution, gently shake it to completely cover the cells, and then incubate at room temperature for 5-30 minutes.

d. Wash twice with medium, 5 minutes each time. Observation by fluorescence microscopy or flow cytometry.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- J Nanobiotechnology. 2023 Sep 2;21(1):311.
- ACS Appl Mater Interfaces. 2023 Dec 14.
- Chem Mater. 2023 Apr 4.
- Int J Biol Macromol. 2023 Oct 4:127219.
- Int J Nanomedicine. 2022, 17: 5229-5246.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Gan WB, et al. Multicolor "DiOlistic" labeling of the nervous system using lipophilic dye combinations. Neuron. 2000 Aug;27(2):219-25.

[2]. Swift MJ, et al. Applied electric fields accelerate the diffusion rate and increase the diffusion distance of DiI in fixed tissue. J Neurosci Methods. 2005 Jan 30;141(1):155-63.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA