



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mag-Fura-2 tetrapotassium

Cat. No.:	HY-D1702
CAS No.:	132319-57-4
Molecular Formula:	C ₁₈ H ₁₀ K ₄ N ₂ O ₁₁
Molecular Weight:	586.67
Target:	Fluorescent Dye
Pathway:	Others
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description	Mag-Fura-2 tetrapotassium is a UV excitable rational fluorescent Mg ²⁺ /Ca ²⁺ indicator (Ex=334-360 nm, Em=510 nm). Mag-Fura-2 tetrapotassium can be used for the determination of Mg ²⁺ and Ca ²⁺ concentrations ^{[1][2]} .
In Vitro	<p>Guidelines (Following is our recommended protocol. This protocol only provides a guideline, and should be modified according to your specific needs).</p> <p>Monitoring of Ca²⁺ release^[1]:</p> <ol style="list-style-type: none"> 1. Incubate dispersed parotid acinar cells with 8 μM Mag-fura-2 tetrapotassium for 30 min at 37°C. (acinar cells for example). 2. Wash cells twice with fresh HBSS-H without BSA. 3. Precoat cell adhesive Cell-Tak on sample chambers. 4. Transfer the dye-loaded cells to the chambers and attach to the bottom. 5. Mount the sample chambers on the stage of an inverted microscope (equipped with a 40 × objective), wash with BSA-free HBSS-H and then with Mg²⁺/ATP-free ICM. 6. Incubate acinar cells with Mg²⁺/ATP-free ICM containing 50 μg/mL saponin for 3-5 min at room temperature. 7. Wash the cells with ICM containing Mg²⁺ and ATP, and incubate in the complete ICM for at least 5 min to allow complete filling of the intracellular Ca²⁺ stores. 8. Alternately excite permeabilised cells, capture and digitise fluorescence emission at 510 nm by a digital imaging system (record the 344 nm/360 nm ratio every 20 s). <p>Note: ICM (intracellular-like medium) containing 125 mM KCl, 19 mM NaCl, 10 mM HEPES (pH 7.3 with KOH), 3 mM ATP, 1.4 mM MgCl₂, 0.33 mM CaCl₂, and 1 mM EGTA.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>

REFERENCES

- [1]. Tojyo Y, et al. Monitoring of Ca²⁺ release from intracellular stores in permeabilized rat parotid acinar cells using the fluorescent indicators Mag-fura-2 and calcium green C18. *Biochem Biophys Res Commun.* 1997 Nov 7;240(1):189-95.
- [2]. Dai LJ, et al. Intracellular Mg²⁺ and magnesium depletion in isolated renal thick ascending limb cells. *J Clin Invest.* 1991 Oct;88(4):1255-64.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA