



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Cystathionine β -lyase, Recombinant Microorganisms

Cat. No.:	HY-E70079		
CAS No.:	9055-05-4		
Target:	Biochemical Assay Reagents		
Pathway:	Others		
Storage:	Powder	-20°C	3 years
	In solvent	-80°C	6 months
		-20°C	1 month

Cystathionine β -lyase, Recombinant Microorganisms

SOLVENT & SOLUBILITY

In Vitro	H ₂ O : 100 mg/mL (Need ultrasonic)
----------	------------------------------------------------

BIOLOGICAL ACTIVITY

Description	Cystathionine β -lyase, Recombinant Microorganisms (CBL) is an enzyme that catalyzes the breakdown of cystathionine to homocysteine, the penultimate step in methionine biosynthesis. Cystathionine β -lyase is important for bacterial virulence ^[1] .
-------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

In Vitro	Cystathionine β -lyase (25 nM; 10 min) catalyzes L-(+)-cystathionine (100 μ M) to homocysteine. Cystathionine β -lyase activity can be measured by monitoring the formation of homocysteine at 412 nm, using 5,5'-dithio-bis-(2-nitrobenzoic acid) as a detector of free thiol ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.
----------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

REFERENCES

[1]. Ejim LJ, et al. Cystathionine beta-lyase is important for virulence of Salmonella enterica serovar Typhimurium. Infect Immun. 2004 Jun;72(6):3310-4.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA