



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Arbutin

Cat. No.:	HY-N0192		
CAS No.:	497-76-7		
Molecular Formula:	C ₁₂ H ₁₆ O ₇		
Molecular Weight:	272.25		
Target:	Tyrosinase; Endogenous Metabolite		
Pathway:	Metabolic Enzyme/Protease		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	2 years
		-20°C	1 year



SOLVENT & SOLUBILITY

In Vitro

DMSO : ≥ 50 mg/mL (183.65 mM)
 H₂O : 33.33 mg/mL (122.42 mM; Need ultrasonic)
 * "≥" means soluble, but saturation unknown.

Preparing Stock Solutions	Solvent		Mass		
	Concentration		1 mg	5 mg	10 mg
	1 mM		3.6731 mL	18.3655 mL	36.7309 mL
	5 mM		0.7346 mL	3.6731 mL	7.3462 mL
	10 mM		0.3673 mL	1.8365 mL	3.6731 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: PBS
Solubility: 100 mg/mL (367.31 mM); Clear solution; Need ultrasonic
- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: ≥ 2.5 mg/mL (9.18 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
Solubility: ≥ 2.5 mg/mL (9.18 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil
Solubility: ≥ 2.5 mg/mL (9.18 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

Arbutin (β-Arbutin) is a competitive inhibitor of tyrosinase, with K_i^{app} values of 1.42 mM for monophenolase; 0.9 mM for diphenolase. Arbutin is also used as depigmenting agents^[1]. Arbutin is a natural polyphenol isolated from the bearberry plant *Arctostaphylos uvaursi*, possesses with anti-oxidant, anti-inflammatory and anti-tumor properties^{[2][3]}.

IC₅₀ & Target	Human Endogenous Metabolite
In Vitro	Arbutin (0.3-5.4 mM; 24 hours, 48 hours, 72 hours; B16 murine melanoma cells) inhibits the viability of B16 murine melanoma cells in a time-and dose-dependent manner ^[2] . ?Arbutin (1.4-5.4 mM; 24 hours) increases the apoptosis rate of B16 murine melanoma cell of treatment at a dose of 5.4 mM [2]. MCE has not independently confirmed the accuracy of these methods. They are for reference only. Cell Viability Assay ^[2]
	Cell Line: B16 murine melanoma cells
	Concentration: 0.3 mM, 0.7 mM, 1.4 mM, 2.1 mM, 2.9 mM, 3.6 mM, 5.4 mM
	Incubation Time: 24 hours, 48 hours or 72 hours
	Result: Inhibited the viability of B16 murine melanoma cells in a time- and dose-dependent manner.
	Apoptosis Analysis ^[2]
	Cell Line: B16 murine melanoma cells
	Concentration: 1.4 mM, 2.9 mM, 5.4 mM
	Incubation Time: 24 hours
	Result: Caused apoptosis in 19.3% of the cells.
In Vivo	Arbutin (50 mg/kg, 100 mg/kg; oral administration; every day; for 17 days; male C57BL/6 mice) pretreatment exhibits markedly protective effects on ISO-induced cardiac hypertrophy in mice ^[3] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.
	Animal Model: Male C57BL/6 mice (20-25 g) ^[3]
	Dosage: 50 mg/kg, 100 mg/kg
	Administration: Oral administration; every day; for 17 days
	Result: Ameliorated the ISO-induced myocardial injury.

CUSTOMER VALIDATION

- FEBS Open Bio. 2021 Jan;11(1):289-299.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Garcia-Jimenez A, et al. Action of tyrosinase on alpha and beta-arbutin: A kinetic study. PLoS One. 2017 May 11;12(5):e0177330.
- [2]. Jiang L, et al. Investigation of the pro-apoptotic effects of arbutin and its acetylated derivative on murinemelanoma cells. Int J Mol Med. 2018 Feb;41(2):1048-1054.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA