



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

L-Tyrosine-¹³C₆

| | | | |
|---------------------------|---|-------|----------|
| Cat. No.: | HY-N0473S2 | | |
| CAS No.: | 201595-63-3 | | |
| Molecular Formula: | C ₃ ¹³ C ₆ H ₁₁ NO ₃ | | |
| Molecular Weight: | 187.14 | | |
| Target: | Endogenous Metabolite | | |
| Pathway: | Metabolic Enzyme/Protease | | |
| Storage: | Powder | -20°C | 3 years |
| | | 4°C | 2 years |
| | In solvent | -80°C | 6 months |
| | | -20°C | 1 month |



SOLVENT & SOLUBILITY

In Vitro

1M HCl : 33.33 mg/mL (178.10 mM; ultrasonic and adjust pH to 1 with HCl)

| Preparing Stock Solutions | Solvent Concentration | Mass | | |
|---------------------------|--------------------------|-----------|------------|------------|
| | | 1 mg | 5 mg | 10 mg |
| | 1 mM | 5.3436 mL | 26.7180 mL | 53.4359 mL |
| | 5 mM | 1.0687 mL | 5.3436 mL | 10.6872 mL |
| | 10 mM | 0.5344 mL | 2.6718 mL | 5.3436 mL |

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

L-Tyrosine-¹³C₆ is the ¹³C-labeled L-Tyrosine. L-Tyrosine is a non-essential amino acid which can inhibit citrate synthase activity in the posterior cortex.

In Vitro

Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. *Ann Pharmacother.* 2019;53(2):211-216.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA