



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Gangliotetraose

Cat. No.:	HY-N10512
CAS No.:	75645-24-8
Molecular Formula:	C <sub>26</sub> H <sub>45</sub> NO <sub>21</sub>
Molecular Weight:	707.63
Target:	Others
Pathway:	Others
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



### BIOLOGICAL ACTIVITY

<b>Description</b>	Gangliotetraose (Gg4) is a tetrasccharide, exhibits major components including GM1 and its sialylated derivatives. GM1 facilitates efflux of nuclear Ca <sup>2+</sup> and reduces the level of nuclear Ca <sup>2+</sup> that characterizes the differentiated neuron. GM1 affects neuronal plasticity and repair mechanisms, as well as neurotrophin release in the brain <sup>[1][2]</sup> .									
<b>IC<sub>50</sub> &amp; Target</b>	Akt, ERK1/2 <sup>[3]</sup> ; amyloid β-protein <sup>[4]</sup>									
<b>In Vitro</b>	<p>Gangliotetraose (GM1) (10 μM; 1 h) increases the viability of pheochromocytoma PC12 cells exposed to hydrogen peroxide (1 mM; 2 h) and diminishes the accumulation of reactive oxygen species and oxidative inactivation of Na<sup>+</sup>, K<sup>+</sup>-ATPase<sup>[3]</sup>.</p> <p>Gangliotetraose (GM1) (100 nM and 10 μM; ) increases the basal activity of Akt and ERK1/2, without changing Akt activity in PC12 cells exposed to hydrogen peroxide<sup>[3]</sup>.</p> <p>Gangliotetraose (GM1) (50 μM; 24 h) binds the midportion of Aβ to produce Aβ oligomers, GM1 bound Aβ (GAβ). GAβ is endogenously generated in the brain and accelerates Aβ assembly by acting as a seed<sup>[4]</sup>.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p> <p>Cell Viability Assay<sup>[3]</sup></p> <table border="1"> <tr> <td>Cell Line:</td> <td>Rat pheochromocytoma PC12 cells</td> </tr> <tr> <td>Concentration:</td> <td>1 nM, 10 nM, 100 nM, 1 μM, 10 μM, 50 μM</td> </tr> <tr> <td>Incubation Time:</td> <td>1 hour (preincubation); started 24 h after the transfer of the cells to the plates; exposed to 1 mM H<sub>2</sub>O<sub>2</sub> for 2 h later</td> </tr> <tr> <td>Result:</td> <td>           Showed protective effect (rescue rates, %) on PC12 cells exposed to H<sub>2</sub>O<sub>2</sub> in a dose-dependent manner.            The rescue rates ranged from 2.7% to 76% with concentration of 1 nM-50 μM.         </td> </tr> </table>		Cell Line:	Rat pheochromocytoma PC12 cells	Concentration:	1 nM, 10 nM, 100 nM, 1 μM, 10 μM, 50 μM	Incubation Time:	1 hour (preincubation); started 24 h after the transfer of the cells to the plates; exposed to 1 mM H <sub>2</sub> O <sub>2</sub> for 2 h later	Result:	Showed protective effect (rescue rates, %) on PC12 cells exposed to H <sub>2</sub> O <sub>2</sub> in a dose-dependent manner. The rescue rates ranged from 2.7% to 76% with concentration of 1 nM-50 μM.
Cell Line:	Rat pheochromocytoma PC12 cells									
Concentration:	1 nM, 10 nM, 100 nM, 1 μM, 10 μM, 50 μM									
Incubation Time:	1 hour (preincubation); started 24 h after the transfer of the cells to the plates; exposed to 1 mM H <sub>2</sub> O <sub>2</sub> for 2 h later									
Result:	Showed protective effect (rescue rates, %) on PC12 cells exposed to H <sub>2</sub> O <sub>2</sub> in a dose-dependent manner. The rescue rates ranged from 2.7% to 76% with concentration of 1 nM-50 μM.									
<b>In Vivo</b>	<p>Gangliotetraose (GM1) (30 mg/kg; i.p.; 5, 11, 42, and 73 d) stimulates the regeneration of nigrostriatal dopaminergic neurons in the central nervous system of rats after unilateral hemitransection<sup>[5]</sup>.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p> <table border="1"> <tr> <td>Animal Model:</td> <td>Unilateral semi-transection model in Sprague-Dawley rats (170-190 g)<sup>[5]</sup></td> </tr> <tr> <td>Dosage:</td> <td>5 mg/kg; 30 mg/kg</td> </tr> </table>		Animal Model:	Unilateral semi-transection model in Sprague-Dawley rats (170-190 g) <sup>[5]</sup>	Dosage:	5 mg/kg; 30 mg/kg				
Animal Model:	Unilateral semi-transection model in Sprague-Dawley rats (170-190 g) <sup>[5]</sup>									
Dosage:	5 mg/kg; 30 mg/kg									

Administration:	Intraperitoneal injection; 5, 11, 42, 73 days; started on day 2 after surgery and finished 24 h before sacrifice
Result:	Increased the $V_{max}$ of tyrosine hydroxylase (TH) in the lesioned side starting on day 14 dose-dependently with 73% (5 mg/kg/d) and 85% (30 mg/kg/d) of that of the unlesioned side, respectively.

## REFERENCES

- [1]. Okada H, et al. Complement-mediated cytolysis and azidothymidine are synergistic in HIV-1 suppression. *Int Immunol*. 1998 Jan;10(1):91-5.
- [2]. Ledeen RW, et al. The role of GM1 and other gangliosides in neuronal differentiation. Overview and new finding. *Ann N Y Acad Sci*. 1998 Jun 19;845:161-75.
- [3]. Zakharova IO, et al. GM1 ganglioside activates ERK1/2 and Akt downstream of Trk tyrosine kinase and protects PC12 cells against hydrogen peroxide toxicity. *Neurochem Res*. 2014 Nov;39(11):2262-75.
- [4]. Toffano G, et al. GM1 ganglioside stimulates the regeneration of dopaminergic neurons in the central nervous system. *Brain Res*. 1983 Feb 14;261(1):163-6.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA