



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

(R)-Reticuline

Cat. No.:	HY-N1356A
CAS No.:	3968-19-2
Molecular Formula:	C ₁₉ H ₂₃ NO ₄
Molecular Weight:	329.39
Target:	Others
Pathway:	Others
Storage:	4°C, protect from light * In solvent : -80°C, 6 months; -20°C, 1 month (protect from light)



SOLVENT & SOLUBILITY

In Vitro	DMSO : 100 mg/mL (303.59 mM; Need ultrasonic)					
	Preparing Stock Solutions	Solvent	Mass	1 mg	5 mg	10 mg
		Concentration				
		1 mM		3.0359 mL	15.1796 mL	30.3591 mL
		5 mM		0.6072 mL	3.0359 mL	6.0718 mL
10 mM		0.3036 mL	1.5180 mL	3.0359 mL		
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (7.59 mM); Clear solution 2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (7.59 mM); Clear solution					

BIOLOGICAL ACTIVITY

Description	<p>(R)-Reticuline is an isomer of Reticuline (HY-N1356). Reticuline displays anti-inflammatory and cardiovascular effects through JAK2/STAT3 and NF-κB signaling pathways. Salutaridine is a key intermediate in morphine biosynthesis. Salutaridine can be converted from (R)-Reticuline in the poppy plant. The conversion system relies on membrane-bound cytochrome P-450 enzymes and also requires reducing cofactors NADPH, molecular oxygen, etc^{[1][2]}.</p>
--------------------	---

REFERENCES

[1]. Gerardy R, et al. Formation of salutaridine from (R)-reticuline by a membrane-bound cytochrome P-450 enzyme from *Papaver somniferum*[J]. *Phytochemistry*, 1992, 32(1): 79-86.

[2]. Yang X, et al. Anti-Inflammatory Effects of Boldine and Reticuline Isolated from *Litsea cubeba* through JAK2/STAT3 and NF- κ B Signaling Pathways. *Planta Med.* 2018 Jan;84(1):20-25.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA