



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

(+)-Coclaurine

Cat. No.:	HY-N2550
CAS No.:	2196-60-3
Molecular Formula:	C ₁₇ H ₁₉ NO ₃
Molecular Weight:	285.34
Target:	Others
Pathway:	Others
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



SOLVENT & SOLUBILITY

In Vitro

DMSO : 25 mg/mL (87.61 mM; ultrasonic and warming and heat to 60°C)

Preparing Stock Solutions	Solvent		1 mg	5 mg	10 mg
	Concentration	Mass			
	1 mM		3.5046 mL	17.5230 mL	35.0459 mL
	5 mM		0.7009 mL	3.5046 mL	7.0092 mL
	10 mM		0.3505 mL	1.7523 mL	3.5046 mL

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

(+)-Coclaurine ((+)-(R)-Coclaurine), benzyltetrahydroisoquinoline alkaloid isolated from a variety of plant sources. (+)-Coclaurine has anti-aging activity^{[1][2]}.

In Vivo

An intracerebroventricular injection of (+)-Coclaurine (d-Coclaurine; 50 µg) produces a slight increase in 3,4-dihydroxyphenylacetic acid level and a significant increase in homovanillic acid level in the mouse striatum. (+)-Coclaurine blocks postsynaptic but not presynaptic dopamine receptors in the mouse striatum^[2].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Siva S Panda, et al. Protective effects of Aporosa octandra bark extract against D-galactose induced cognitive impairment and oxidative stress in mice. Heliyon. 2018 Nov 30;4(11):e00951.

[2]. H Watanabe, et al. Effects of d-coclaurine and d-reticuline, benzyltetrahydroisoquinoline alkaloids, on levels of 3,4-dihydroxyphenylacetic acid and homovanillic acid in the mouse striatum. J Pharmacobiodyn. 1983 Oct;6(10):793-6.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA