



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Sterigmatocystine-¹³C₁₈

Cat. No.:	HY-N6725S
Molecular Formula:	¹³ C ₁₈ H ₁₂ O ₆
Molecular Weight:	342.15
Target:	Endogenous Metabolite; Apoptosis; Antibiotic; Bacterial; DNA/RNA Synthesis; Isotope-Labeled Compounds
Pathway:	Metabolic Enzyme/Protease; Apoptosis; Anti-infection; Cell Cycle/DNA Damage; Others
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description	Sterigmatocystine- ¹³ C ₁₈ is ¹³ C labeled 2,6-Dimethoxyphenol (HY-W003972). 2,6-Dimethoxyphenol is a phenolic compound that is extensively used for the measurement of laccase activity ^[1] .
In Vitro	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs ^[1] . Sterigmatocystine-induced DNA damage activates the ATM/53-dependent signaling pathway, which contributes to the induction of G2 arrest in GES-1 cells ^[5] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.
In Vivo	Sterigmatocystine (ip; 3 mg/kg once daily for 14 days) inhibits p21 ^{WAF1/CIP1} ^[4] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

- [1]. Kusunoki M, et al. Long-term administration of the fungus toxin, sterigmatocystin, induces intestinal metaplasia and increases the proliferative activity of PCNA, p53, and MDM2 in the gastric mucosa of aged Mongolian gerbils. *Environ Health Prev Med.* 2011 Jul;16(4):224-31.
- [2]. Tong YF, et al. Cyclin-Dependent Kinase Inhibitor p21WAF1/CIP1 Facilitates the Development of Cardiac Hypertroph. *Cell Physiol Biochem.* 2017;42(4):1645-1656.
- [3]. Schroeder HW, et al. Production of sterigmatocystin by some species of the genus *Aspergillus* and its toxicity to chicken embryos. *Appl Microbiol.* 1975 Oct;30(4):589-91.
- [4]. Zhang D, et al. Sterigmatocystin-induced DNA damage triggers G2 arrest via an ATM/p53-related pathway in human gastric epithelium GES-1 cells in vitro. *PLoS One.* 2013 May 21;8(5):e65044.
- [5]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. *Ann Pharmacother.* 2019 Feb;53(2):211-216.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA