



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Desoxo-narchinol A

Cat. No.:	HY-N8435
CAS No.:	53859-06-6
Molecular Formula:	C ₁₂ H ₁₆ O ₂
Molecular Weight:	192.25
Target:	TNF Receptor; COX; NO Synthase; PGE synthase
Pathway:	Apoptosis; Immunology/Inflammation
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description	Desoxo-narchinol A is an orally active and potent anti-inflammatory agent. Desoxo-narchinol A can be isolated from the roots and rhizomes of <i>Nardostachys jatamansi</i> . Desoxo-narchinol A can be used for septic shock and inflammatory diseases research ^{[1][2][3]} .
In Vitro	Desoxo-narchinol A inhibits tissue injury and production of pro-inflammatory cytokines, such as IL-1 β , IL-6, and TNF- α , in the liver and lung ^[2] . Desoxo-narchinol A (0-500 nM, 24 h) inhibits the production of inflammatory mediators, such as iNOS and its derivative NO, COX-2, PGE2, IL-1 β , IL-6 and TNF- α and H3 protein acetylation in murine peritoneal macrophages ^[2] . Desoxo-narchinol A inhibits LPS-induced activation of p38 in murine macrophages ^[2] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.
In Vivo	Desoxo-narchinol A (0-0.5 mg/kg, IP, once) dramatically reduced mortality in a murine LPS-induced endotoxin shock model ^[2] . Desoxo-narchinol A (50 mg/kg, PO, once) shows the oral bioavailability of 18.1% in rats and 28.4% in mice ^[3] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

- [1]. Itokawa H, et al. Cytotoxic sesquiterpenes from *Nardostachys chinensis*. *Chem Pharm Bull (Tokyo)*. 1993 Jun;41(6):1183-4.
- [2]. Shin JY, et al. Anti-inflammatory effect of desoxo-narchinol-A isolated from *Nardostachys jatamansi* against lipopolysaccharide. *Int Immunopharmacol*. 2015 Dec;29(2):730-738.
- [3]. Thapa SK, Upadhyay M, Kim TH, Shin S, Park SJ, Shin BS. Liquid Chromatography-Tandem Mass Spectrometry of Desoxo-Narchinol a and Its Pharmacokinetics and Oral Bioavailability in Rats and Mice. *Molecules*. 2019 May 28;24(11):2037.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA