



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



## Malabaricone B

Cat. No.: HY-N8517

CAS No.: 63335-24-0

Molecular Formula: C<sub>21</sub>H<sub>26</sub>O<sub>4</sub>


Molecular Weight: 342.43

Target: Glucosidase; Apoptosis

Pathway: Metabolic Enzyme/Protease; Apoptosis

Storage: 4°C, protect from light

\* In solvent : -80°C, 6 months; -20°C, 1 month (protect from light)



### SOLVENT & SOLUBILITY

#### In Vitro

DMSO : 100 mg/mL (292.03 mM; Need ultrasonic)

Preparing Stock Solutions	Concentration	Mass		
		1 mg	5 mg	10 mg
	1 mM	2.9203 mL	14.6015 mL	29.2030 mL
	5 mM	0.5841 mL	2.9203 mL	5.8406 mL
	10 mM	0.2920 mL	1.4602 mL	2.9203 mL

Please refer to the solubility information to select the appropriate solvent.

### BIOLOGICAL ACTIVITY

#### Description

Malabaricone B, a naturally occurring plant phenolic, is an orally active  $\alpha$ -glucosidase inhibitor with an IC<sub>50</sub> of 63.7  $\mu$ M. Malabaricone B has anticancer, antimicrobial, anti-oxidation and antidiabetic activities<sup>[1][2][3]</sup>.

#### In Vitro

Malabaricone B shows selective toxicity to human lung cancer (A549), malignant melanoma (A375) and T cell leukemia (Jurkat) cell lines, without showing toxicity to human normal intestinal (INT407), human kidney (HEK293) and lung fibroblast (WI-38) cells. Among the chosen cancer cell lines, Malabaricone B shows maximum cytotoxicity to the A549 cells (IC<sub>50</sub> = 8.1  $\mu$ M), which is significantly better than that of curcumin (IC<sub>50</sub> = 26.7  $\mu$ M)<sup>[1]</sup>.

The Malabaricone B-induced apoptosis is mediated by an increase in the intracellular reactive oxygen species (ROS). Malabaricone B (2.5, 5, 10 and 20  $\mu$ M) increases the BAX level while simultaneously decreasing the BCL-2 and BCL-XL levels in the A549 cells, triggering the mitochondrial apoptotic pathway as revealed from the release of cytochrome c, and the activation of caspase-9 and caspase-3<sup>[1]</sup>.

Malabaricone B exhibits a good level of antimicrobial activity when tested against a variety of microorganisms, including *Staphylococcus aureus* and *Candida albicans*<sup>[2]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### In Vivo

Malabaricone B (100 mg/kg; p.o; alternate day, 23 days) inhibits lung tumor (xenograft) growth in SCID mice<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

---

## REFERENCES

---

- [1]. Mrityunjay Tyagi, et al. Spice-derived phenolic, malabaricone B induces mitochondrial damage in lung cancer cells via a p53-independent pathway. *Food Funct.* 2018 Nov 14;9(11):5715-5727.
- [2]. K Y Orabi, et al. Isolation and characterization of two antimicrobial agents from mace (*Myristica fragrans*). *J Nat Prod.* May-Jun 1991;54(3):856-9.
- [3]. B Prabha, et al. Antidiabetic potential of phytochemicals isolated from the stem bark of *Myristica fatua* Houtt. var. *magnifica* (Bedd.) Sinclair. *Bioorg Med Chem.* 2018 Jul 23;26(12):3461-3467.
- 

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA