



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Suc-Leu-Leu-Val-Tyr-AMC

<b>Cat. No.:</b>	HY-P1002		
<b>CAS No.:</b>	94367-21-2		
<b>Molecular Formula:</b>	C <sub>40</sub> H <sub>53</sub> N <sub>5</sub> O <sub>10</sub>		
<b>Molecular Weight:</b>	763.88		
<b>Sequence:</b>	Suc-Leu-Leu-Val-Tyr		
<b>Sequence Shortening:</b>	Suc-LLVY		
<b>Target:</b>	Fluorescent Dye		
<b>Pathway:</b>	Others		
<b>Storage:</b>	Pure form	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



### SOLVENT & SOLUBILITY

#### In Vitro

DMSO :  $\geq 20$  mg/mL (26.18 mM)  
 \* " $\geq$ " means soluble, but saturation unknown.

	Solvent Concentration	Mass		
		1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM	1.3091 mL	6.5455 mL	13.0911 mL
	5 mM	0.2618 mL	1.3091 mL	2.6182 mL
	10 mM	0.1309 mL	0.6546 mL	1.3091 mL

Please refer to the solubility information to select the appropriate solvent.

### BIOLOGICAL ACTIVITY

#### Description

Suc-Leu-Leu-Val-Tyr-AMC is a fluorogenic substrate.

#### In Vitro

Suc-Leu-Leu-Val-Tyr-AMC (Suc-LLVY) is a membrane-permeable calpain-specific fluorogenic substrate, proteolytic hydrolysis of the peptidyl-7-amino bond liberates the highly fluorescent 7-amino-4-methylcoumarin (AMC) moiety<sup>[1]</sup>. The effect of TGF- $\beta$  on hydrolysis of these substrates (e.g Suc-Leu-Leu-Val-Tyr-AMC) are assessed. Biliary epithelial H69 cells are incubated with 10, 1, 0.1, or 0 ng/mL TGF- $\beta$  for 24 h. Substrate hydrolysis is then fluorometrically assessed in cytosolic extracts. Basal activity is 1.12, 8.33, and 14.52 nmol AMC/mg protein/min for suc-LLVY-AMC, z-LLE-AMC, and z-LLL-AMC hydrolysis, respectively<sup>[2]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## PROTOCOL

### Kinase Assay <sup>[3]</sup>

Immunoprecipitation is carried out for the two sets of samples, using the same amount of protein. The 20 S and 26 S proteasome immunoprecipitates are washed with 50 mM Hepes/KOH (pH 7.5), and 50 mM Hepes/KOH (pH 7.5) containing 2 mM ATP, respectively, prior to the determination of peptidase activity using 50  $\mu$ M suc-Leu- Leu-Val-Tyr-AMC as substrate in these buffers<sup>[3]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## CUSTOMER VALIDATION

- Nature. 2022 May;605(7910):567-574.
- Environ Sci Technol. 2019 Aug 20;53(16):9789-9799.
- Food Funct. 20 Sep 2021.
- BMC Dev Biol. 2021 Feb 1;21(1):4.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

[1]. Roberta L. DeBiasi, et al. Reovirus-Induced Apoptosis Is Preceded by Increased Cellular Calpain Activity and Is Blocked by Calpain Inhibitors. J Virol. 1999 Jan; 73(1): 695–701.

[2]. Tadlock L, et al. Transforming growth factor-beta inhibition of proteasomal activity: a potential mechanism of growth arrest. Am J Physiol Cell Physiol. 2003 Aug;285(2):C277-85. Epub 2003 Mar 19.

[3]. Gardner RC, et al. Characterization of peptidyl boronic acid inhibitors of mammalian 20 S and 26 S proteasomes and their inhibition of proteasomes in cultured cells. Biochem J. 2000 Mar, 2:447-54.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA