



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

VIP(Guinea pig)

Cat. No.:	HY-P1015
CAS No.:	96886-24-7
Molecular Formula:	C ₁₄₇ H ₂₃₉ N ₄₃ O ₄₂ S ₂
Molecular Weight:	3344.86
Sequence:	His-Ser-Asp-Ala-Leu-Phe-Thr-Asp-Thr-Tyr-Thr-Arg-Leu-Arg-Lys-Gln-Met-Ala-Met-Lys-Lys-Tyr-Leu-Asn-Ser-Val-Leu-Asn-NH ₂ HSDALFTDTYTRLRKQMAMKKYLNSVLN-NH ₂
Sequence Shortening:	HSDALFTDTYTRLRKQMAMKKYLNSVLN-NH ₂
Target:	Others
Pathway:	Others
Storage:	Sealed storage, away from moisture and light, under nitrogen Powder -80°C 2 years -20°C 1 year * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light, under nitrogen)

SOLVENT & SOLUBILITY

In Vitro	H ₂ O : ≥ 1 mg/mL (0.30 mM) * "≥" means soluble, but saturation unknown.
-----------------	--

BIOLOGICAL ACTIVITY

Description	VIP Guinea pig (Vasoactive intestinal peptide), a trophic and mitogenic factor, stimulates growth in whole cultured embryos. VIP Guinea pig functions as a simple gastrointestinal hormone and suggest a possible neurotransmitter function ^{[1][2]} .
In Vitro	Vasoactive intestinal peptide (10 ⁻⁷ M) shortened S phase and G1 phase of neuroepithelial cells by 50% (4.8-2.4 h) and 58% (1.9-0.8 h), respectively, compared with controls ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

- [1]. P Gressens, et al. Vasoactive intestinal peptide shortens both G1 and S phases of neural cell cycle in whole postimplantation cultured mouse embryos. Eur J Neurosci. 1998 May;10(5):1734-42.
- [2]. M G Bryant, et al. Possible dual role for vasoactive intestinal peptide as gastrointestinal hormone and neurotransmitter substance. Lancet. 1976 May 8;1(7967):991-3.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA