



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Thermolysin, *Bacillus thermoproteolyticus* rokko

Cat. No.:	HY-P1748
CAS No.:	9073-78-3
Target:	Endogenous Metabolite
Pathway:	Metabolic Enzyme/Protease
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.

Thermolysin

SOLVENT & SOLUBILITY

In Vitro	H ₂ O : 1.96 mg/mL (adjust pH to 3 with HCl)
----------	---

BIOLOGICAL ACTIVITY

Description Thermolysin, *Bacillus thermoproteolyticus* rokko (EC 3.4.24.27) (TML) is a thermostable neutral metalloproteinase enzyme secreted by the Gram-positive bacteria *Bacillus thermoproteolyticus*. Thermolysin catalyzes the hydrolysis of peptide bonds containing hydrophobic residues^[1].

Optimal pH: 8.0. Considerably stable from pH 5 to 9.5.
Optimal temperature : 70 °C

In Vitro Thermolysin (100 µg in 1 mL, 64°C for 4 h, in isolated starch granules) digestion results in the selective removal of a group of low-molecular-mass proteins ranging between 10 and 27 kD^[2]. Thermolysin can be used for the hydrolysis of bovine liver sarcoplasmic proteins^[3]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo Thermolysin (50 mg/kg, p.o., once) lowers systolic blood pressure in SHR rats^[4]. Thermolysin is not acutely toxic with an oral LD₅₀ of more than 18000 mg/kg in rats and more than 24000 mg/kg in mice (Acute toxicity studies)^[5]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	Spontaneously hypertensive rats (SHR) rats ^[4]
Dosage:	50 mg/kg
Administration:	Oral administration (p.o.), once
Result:	Lowered systolic blood pressure in SHR rats, with maximal reduction by 22 mm in systolic blood pressure being observed 6 h after administration.

REFERENCES

- [1]. Bertusvan den Burg, et al. Chapter 111 - Thermolysin and Related Bacillus Metallopeptidases.
- [2]. Mu-Forster C, et al. Surface localization of zein storage proteins in starch granules from maize endosperm. Proteolytic removal by thermolysin and in vitro cross-linking of granule-associated polypeptides. *Plant Physiol.* 1998 Apr;116(4):1563-71.
- [3]. Di Bernardini R, et al. Isolation, purification and characterization of antioxidant peptidic fractions from a bovine liver sarcoplasmic protein thermolysin hydrolyzate. *Peptides.* 2011 Feb;32(2):388-400.
- [4]. Fujita H, et al: a prodrug-type ACE-inhibitory peptide derived from fish protein. *Immunopharmacology.* 1999 Oct 15;44(1-2):123-7.
- [5]. Ke Q, et al. Safety evaluation of a thermolysin enzyme produced from *Geobacillus stearothermophilus*. *Food Chem Toxicol.* 2013 Sep;59:541-8.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA