



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Phalloidin-TRITC

Cat. No.:	HY-P2270
CAS No.:	915013-10-4
Molecular Formula:	$C_{60}H_{70}N_{12}O_{13}S_2$
Molecular Weight:	1231.4
Target:	Arp2/3 Complex; Fluorescent Dye
Pathway:	Cytoskeleton; Others
Storage:	Sealed storage, away from moisture and light, under nitrogen Powder -80°C 2 years -20°C 1 year * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light, under nitrogen)

BIOLOGICAL ACTIVITY

Description	Phalloidin-TRITC is a fluorescein derivative of Phalloidin, which can specifically label myofilin and display red fluorescence when labeled and can be observed using Tetra channels ^[1] .
In Vitro	<ol style="list-style-type: none"> 1. Preparation of Phalloidin-TRITC working solution <ol style="list-style-type: none"> 1.1 Preparation of the stock solution Dissolve Phalloidin-TRITC in Methanol to obtain 10 mM of stock solution. Note: It is recommended to store the stock solution at -20 °C or -80 °C away from light and avoid repetitive freeze-thaw cycles. 1.2 Preparation of Phalloidin-TRITC working solution Dilute the stock solution in serum-free cell culture medium to obtain 1-10 μM of working solution. Note: Please adjust the concentration of Phalloidin-TRITC working solution according to the actual situation. 2. Cell staining <ol style="list-style-type: none"> 2.1 Suspension cells (6-well plate) <ol style="list-style-type: none"> a. Centrifuge at 1000 g at 4°C for 3-5 minutes and then discard the supernatant. Wash twice with PBS, 5 minutes each time. The cell density is 1×10^6/mL. b. Add 1 mL of working solution, and then incubate at room temperature for 30-60 minutes. c. Centrifuge at 400 g at 4°C for 3-4 minutes and then discard the supernatant. d. Wash twice with PBS, 5 minutes each time. e. Resuspend cells with serum-free cell culture medium or PBS. Observation by fluorescence microscopy or flow cytometry. 2.2 Adherent cells <ol style="list-style-type: none"> a. Culture adherent cells on sterile coverslips. b. Remove the coverslip from the medium and aspirate excess medium. c. Add 100 μL of working solution, gently shake it to completely cover the cells, and then incubate at room temperature for 30-60 minutes. d. Wash twice with medium, 5 minutes each time. Observation by fluorescence microscopy. <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>

CUSTOMER VALIDATION

-
- Small. 2022 Jun 9;e2201147.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. J A Cooper, et al. Effects of cytochalasin and phalloidin on actin. J Cell Biol. 1987 Oct;105(4):1473-8.

[2]. J Wehland, et al. Phalloidin-induced actin polymerization in the cytoplasm of cultured cells interferes with cell locomotion and growth. Proc Natl Acad Sci U S A. 1977 Dec;74(12):5613-7.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA