



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



2,4,6-Trihydroxybenzaldehyde

Cat. No.: HY-W005130

CAS No.: 487-70-7

Molecular Formula: C₇H₆O₄


Molecular Weight: 154.12

Target: NF-κB

Pathway: NF-κB

Storage: 4°C, stored under nitrogen

* In solvent : -80°C, 6 months; -20°C, 1 month (stored under nitrogen)



BIOLOGICAL ACTIVITY

Description	2,4,6-Trihydroxybenzaldehyde is an orally active NF-κB inhibitor. 2,4,6-Trihydroxybenzaldehyde shows anti-tumor activity, anti-cancer cell proliferative activity and anti-obesity activity ^{[1][2][3]} .
In Vitro	2,4,6-Trihydroxybenzaldehyde inhibits adipocyte differentiation and lipid accumulation in 3T3-L1 cells ^[1] . 2,4,6-Trihydroxybenzaldehyde down-regulates PPAR γ , C/EBP α , SREBP-1c, and FAS protein expression levels ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.
In Vivo	2,4,6-Trihydroxybenzaldehyde (oral administration; 5 and 25 mg/kg for 13 weeks) reduces the HFD-induced increase in weight gain, reduces serum levels of glucose, triglycerides, and total cholesterol ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

- [1]. Kim KN, et al. 2,4,6-Trihydroxybenzaldehyde, a potential anti-obesity treatment, suppressed adipocyte differentiation in 3T3-L1 cells and fat accumulation induced by high-fat diet in C57BL/6 mice. Environ Toxicol Pharmacol. 2015 Mar;39(2):962-8.
- [2]. Marton A, et al. Vanillin Analogues o-Vanillin and 2,4,6-Trihydroxybenzaldehyde Inhibit NF κ B Activation and Suppress Growth of A375 Human Melanoma. Anticancer Res. 2016 Nov;36(11):5743-5750.
- [3]. Forester SC, et al. Gut metabolites of anthocyanins, gallic acid, 3-O-methylgallic acid, and 2,4,6-trihydroxybenzaldehyde, inhibit cell proliferation of Caco-2 cells. J Agric Food Chem. 2010 May 12;58(9):5320-7.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA