



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Fmoc-2-Nal-OH

<b>Cat. No.:</b>	HY-W010893		
<b>CAS No.:</b>	112883-43-9		
<b>Molecular Formula:</b>	C <sub>28</sub> H <sub>23</sub> NO <sub>4</sub>		
<b>Molecular Weight:</b>	437.49		
<b>Target:</b>	Amino Acid Derivatives		
<b>Pathway:</b>	Others		
<b>Storage:</b>	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



### SOLVENT & SOLUBILITY

<b>In Vitro</b>	DMSO : 100 mg/mL (228.58 mM; Need ultrasonic)			
		<b>Solvent</b>	<b>Mass</b>	
		<b>Concentration</b>	<b>1 mg</b>	<b>5 mg</b>
	<b>Preparing Stock Solutions</b>		<b>10 mg</b>	
		<b>1 mM</b>	2.2858 mL	11.4288 mL
		<b>5 mM</b>	0.4572 mL	2.2858 mL
		<b>10 mM</b>	0.2286 mL	1.1429 mL
	Please refer to the solubility information to select the appropriate solvent.			
<b>In Vivo</b>	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2 mg/mL (4.57 mM); Clear solution  2. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2 mg/mL (4.57 mM); Clear solution			

### BIOLOGICAL ACTIVITY

<b>Description</b>	Fmoc-2-Nal-OH is an alanine derivative <sup>[1]</sup> .
<b>In Vitro</b>	Amino acids and amino acid derivatives have been commercially used as ergogenic supplements. They influence the secretion of anabolic hormones, supply of fuel during exercise, mental performance during stress related tasks and prevent exercise induced muscle damage. They are recognized to be beneficial as ergogenic dietary substances <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### REFERENCES

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA