



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## 3-Methoxyphenylacetic acid

Cat. No.:	HY-W015343		
CAS No.:	1798-09-0		
Molecular Formula:	C <sub>9</sub> H <sub>10</sub> O <sub>3</sub>		
Molecular Weight:	166.17		
Target:	Endogenous Metabolite		
Pathway:	Metabolic Enzyme/Protease		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



### SOLVENT & SOLUBILITY

In Vitro	DMSO : 100 mg/mL (601.79 mM; Need ultrasonic)			
		Solvent Concentration	Mass	
			1 mg	5 mg
	Preparing Stock Solutions		10 mg	
	1 mM	6.0179 mL	30.0897 mL	60.1793 mL
	5 mM	1.2036 mL	6.0179 mL	12.0359 mL
	10 mM	0.6018 mL	3.0090 mL	6.0179 mL
Please refer to the solubility information to select the appropriate solvent.				
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (15.04 mM); Clear solution			
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (15.04 mM); Clear solution			
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (15.04 mM); Clear solution			

### BIOLOGICAL ACTIVITY

Description	3-Methoxyphenylacetic acid (m-Methoxyphenylacetic acid), a m-hydroxyphenylacetic acid (m-OHPAA) derivative, is a phytotoxin in <i>Rhizoctonia solani</i> . 3-Methoxyphenylacetic acid is used to develop a toxin-mediated bioassay for resistance to <i>rhizoctonia root rot</i> <sup>[1]</sup> .
-------------	---

### REFERENCES

---

[1]. Mandava NB, et al. Phytotoxins in *Rhizoctonia solani*: isolation and biological activity of m-hydroxy- and m-methoxyphenylacetic acids. *J Agric Food Chem*. 1980 Jan-Feb;28(1):71-5.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA