



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Sodium 2-oxopropanoate

<b>Cat. No.:</b>	HY-W015913
<b>CAS No.:</b>	113-24-6
<b>Molecular Formula:</b>	C <sub>3</sub> H <sub>3</sub> NaO <sub>3</sub>
<b>Molecular Weight:</b>	110
<b>Target:</b>	Endogenous Metabolite; Reactive Oxygen Species
<b>Pathway:</b>	Metabolic Enzyme/Protease; Immunology/Inflammation; NF-κB
<b>Storage:</b>	4°C, sealed storage, away from moisture * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture)



### SOLVENT & SOLUBILITY

<b>In Vitro</b>	H <sub>2</sub> O : 100 mg/mL (909.09 mM; Need ultrasonic)				
		Solvent Concentration	Mass		
	<b>Preparing Stock Solutions</b>		1 mg	5 mg	10 mg
		1 mM	9.0909 mL	45.4545 mL	90.9091 mL
		5 mM	1.8182 mL	9.0909 mL	18.1818 mL
	10 mM	0.9091 mL	4.5455 mL	9.0909 mL	
Please refer to the solubility information to select the appropriate solvent.					
<b>In Vivo</b>	1. Add each solvent one by one: PBS Solubility: 50 mg/mL (454.55 mM); Clear solution; Need ultrasonic				

### BIOLOGICAL ACTIVITY

<b>Description</b>	Sodium 2-oxopropanoate (Sodium pyruvate), a three-carbon metabolite of Glucose, is a compound produced in the glycolytic pathway. Sodium 2-oxopropanoate is a free radical scavenger that can scavenge ROS <sup>[1][2]</sup> .	
<b>IC<sub>50</sub> &amp; Target</b>	Microbial Metabolite	Human Endogenous Metabolite
<b>In Vitro</b>	<p>In the process of scavenging hydrogen peroxide, Sodium 2-oxopropanoate (Sodium pyruvate) is decarboxylated to lactate and hence diverted away from aerobic metabolism and further ROS generation. Under conditions in which pyruvate stimulated JNK1 activity, both mitochondrial and cytosolic ROS levels rose. An increase in JNK1 activity in a variety of different cell types over a range of pyruvate concentrations are observed<sup>[1]</sup>.</p> <p>Sodium 2-oxopropanoate (Sodium pyruvate) is an effective scavenger of H<sub>2</sub>O<sub>2</sub> as well as of O<sub>2</sub><sup>-</sup>, thereby protecting the lens against oxidative stress and consequent cataract formation, under in vitro as well as in vivo conditions, Sodium 2-oxopropanoate has also been shown to protect the lens proteins against glycation by competitively inhibiting the initial reaction between the sugar carbonyl and the protein -NH<sub>2</sub><sup>[2]</sup>.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>	

## In Vivo

Neonatal rats are given a bolus of Sodium 2-oxopropanoate (Sodium pyruvate; 0.1-10 g/kg), and 1 h later, levels of JNK1 activity were determined in liver extracts. Sodium 2-oxopropanoate (Sodium pyruvate) addition resulted in a significant increase in JNK1 activity. Analysis of serum levels of Sodium 2-oxopropanoate in neonatal rats revealed that baseline levels of pyruvate are 0.30 mM. At the time of sacrifice, following the largest dose of Sodium 2-oxopropanoate, serum levels rose approximately sixfold to 1.84 mM<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## CUSTOMER VALIDATION

- Cell Death Dis. 2023 Apr 6;14(4):246.
- Cell Prolif. 2023 Apr 21;e13442.
- Int Immunopharmacol. 2023 May 12;120:110292.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

[1]. S Nemoto, et al. Role for mitochondrial oxidants as regulators of cellular metabolism. Mol Cell Biol. 2000 Oct;20(19):7311-8.

[2]. W Zhao, et al. Fructose induced deactivation of antioxidant enzymes: preventive effect of pyruvate. Free Radic Res. 2000 Jul;33(1):23-30.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA