



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Acivicin

Cat. No.:	HY-W016586
CAS No.:	42228-92-2
Molecular Formula:	C <sub>5</sub> H <sub>7</sub> ClN <sub>2</sub> O <sub>3</sub>
Molecular Weight:	178.57
Target:	Parasite
Pathway:	Anti-infection
Storage:	4°C, stored under nitrogen * In solvent : -80°C, 6 months; -20°C, 1 month (stored under nitrogen)



### SOLVENT & SOLUBILITY

In Vitro H<sub>2</sub>O : 6.67 mg/mL (37.35 mM; Need ultrasonic)

Preparing Stock Solutions	Solvent Concentration	Mass		
		1 mg	5 mg	10 mg
	1 mM	5.6000 mL	28.0002 mL	56.0004 mL
	5 mM	1.1200 mL	5.6000 mL	11.2001 mL
	10 mM	0.5600 mL	2.8000 mL	5.6000 mL

Please refer to the solubility information to select the appropriate solvent.

### BIOLOGICAL ACTIVITY

**Description** Acivicin (AT-125), a natural product produced by *Streptomyces sviveus* is a  $\gamma$ -glutamyl transpeptidase (GGT) inhibitor. Acivicin can cross the blood-brain barrier and has anti-cancer, anti-parasitic properties<sup>[1][2]</sup>.

**IC<sub>50</sub> & Target**  $\gamma$ -glutamyl transpeptidase<sup>[1]</sup>

**In Vitro** Acivicin (AT-125; 0.1-50  $\mu$ M; 5 days) has an IC<sub>50</sub> of 0.7  $\mu$ M in human HepG2 cells<sup>[1]</sup>.  
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

**In Vivo** Acivicin (AT-125; 5 mg/kg; IP; twice weekly) reduces urinary  $\gamma$ -GT by 70-78%<sup>[3]</sup>.  
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	Male pigmented Long-Evans rats weighed between 250 g and 300 g exposed to Toluene <sup>[3]</sup>
Dosage:	5 mg/kg
Administration:	IP; twice weekly (monday and wednesday)

---

Result:	Reduced urinary $\gamma$ -GT by 70-78%.
---------	---

---

## REFERENCES

- [1]. Kreuzer J, et al. Target discovery of acivicin in cancer cells elucidates its mechanism of growth inhibition. Chem Sci. 2014 Dec 1;6(1):237-245. Epub 2014 Sep 26.
- [2]. Chikhale EG, et al. Carrier-mediated transport of the antitumor agent acivicin across the blood-brain barrier. Biochem Pharmacol. 1995 Mar 30;49(7):941-5.
- [3]. Delphine Waniusiow, et al. Toluene-induced hearing loss in acivicin-treated rats. Neurotoxicol Teratol. May-Jun 2008;30(3):154-60.
- 

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA