



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

Xanthine

Cat. No.:	HY-W017389		
CAS No.:	69-89-6		
Molecular Formula:	C ₅ H ₄ N ₄ O ₂		
Molecular Weight:	152.11		
Target:	Endogenous Metabolite		
Pathway:	Metabolic Enzyme/Protease		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro

1M NaOH : 8.33 mg/mL (54.76 mM; ultrasonic and adjust pH to 11 with NaOH)
 H₂O : 5 mg/mL (32.87 mM; ultrasonic and warming and adjust pH to 10 with 1M NaOH and heat to 60°C)
 DMSO : 3.33 mg/mL (21.89 mM; ultrasonic and warming and heat to 60°C)

Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg
		Concentration	1 mg	5 mg	10 mg
	1 mM		6.5742 mL	32.8709 mL	65.7419 mL
	5 mM		1.3148 mL	6.5742 mL	13.1484 mL
	10 mM		0.6574 mL	3.2871 mL	6.5742 mL

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

Xanthine, a plant alkaloid found in tea, coffee, and cocoa, is a mild stimulant of the central nervous system. Xanthine also acts as an intermediate product on the pathway of purine degradation^{[1][2][3]}.

IC₅₀ & Target

Microbial Metabolite Human Endogenous Metabolite

In Vitro

A number of stimulants are derived from Xanthine including caffeine and theobromine. Xanthine is a product on the pathway of purine degradation. Xanthine is subsequently converted to uric acid by the action of the Xanthine oxidase enzyme.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Talanta. 2023 Sep 6, 125171.
- Research Square Preprint. 2022 Feb.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Pundir CS, Devi R. Biosensing methods for xanthine determination: a review. Enzyme Microb Technol. 2014;57:55-62.
- [2]. David A. Taylor, et al. Central Nervous System Stimulants.
- [3]. Ashihara H, et al. Xanthine Alkaloids: Occurrence, Biosynthesis, and Function in Plants. Prog Chem Org Nat Prod. 2017;105:1-88.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA