



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## MCA

Cat. No.:	HY-W027544
CAS No.:	62935-72-2
Molecular Formula:	C <sub>12</sub> H <sub>10</sub> O <sub>5</sub>
Molecular Weight:	234.2
Target:	Fluorescent Dye
Pathway:	Others
Storage:	4°C, protect from light * In solvent : -80°C, 6 months; -20°C, 1 month (protect from light)



## SOLVENT & SOLUBILITY

In Vitro	DMSO : 25 mg/mL (106.75 mM; Need ultrasonic)				
		Solvent Concentration	Mass		
	Preparing Stock Solutions		1 mg	5 mg	10 mg
		1 mM	4.2699 mL	21.3493 mL	42.6985 mL
		5 mM	0.8540 mL	4.2699 mL	8.5397 mL
	10 mM	0.4270 mL	2.1349 mL	4.2699 mL	
Please refer to the solubility information to select the appropriate solvent.					
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (10.67 mM); Clear solution				
	2. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (10.67 mM); Clear solution				

## BIOLOGICAL ACTIVITY

Description	MCA is a coumarin derivative. MCA quantitates platelet-activating factor by high-performance liquid chromatography with fluorescent detection. MCA can modify FRET peptide substrates for analyzing protease activities <sup>[1][2]</sup> .
-------------	---

## REFERENCES

- [1]. Perera NC, et, al. NSP4 is stored in azurophil granules and released by activated neutrophils as active endoprotease with restricted specificity. J Immunol. 2013 Sep 1;191(5):2700-7.
- [2]. Choi YH, et, al. Effect of functional groups on the solubilities of coumarin derivatives in supercritical carbon dioxide. Chromatographia. 1998 Jan;47(93-7).

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA