



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Dehydrothio-p-toluidine

| | |
|--------------------|--|
| Cat. No.: | HY-W040269 |
| CAS No.: | 92-36-4 |
| Molecular Formula: | C ₁₄ H ₁₂ N ₂ S |
| Molecular Weight: | 240.32 |
| Target: | Fluorescent Dye |
| Pathway: | Others |
| Storage: | 4°C, protect from light * In solvent : -80°C, 6 months; -20°C, 1 month (protect from light) |



BIOLOGICAL ACTIVITY

| | |
|--------------------|--|
| Description | Dehydrothio-p-toluidine (DHPT) is a chemical intermediate used in dyestuff production. Dehydrothio-p-toluidine shows inhalation toxicity in acute and subchronic conditions ^[1] . |
| In Vivo | <p>Dehydrothio-p-toluidine (4 h) induces salivation, pawing and chewing motions, lachrymation, sporadic fasciculations, rapid respiration and slight weight loss of mice under an average of 0.64 mg DHPT/L condition^[1].</p> <p>Dehydrothio-p-toluidine (4 h) induces sporadic nasal discharge and lethargy of mice under an average of 2.41 mg DHPT/L condition^[1].</p> <p>Dehydrothio-p-toluidine (6 h/day on 5 days followed by two rest days and then by five additional days of exposures; 0.6 mg DHPT/L) decreases the erythrocyte count, haemoglobin level and relative number of lymphocytes than in the controls, excretes a larger volume of less-concentrated urine, produces liver changes characterized by hepatocyte hyper trophy and proliferation of bile-duct epithelial cells and induces thymic necrosis and depletion of lymphocytes after the tenth exposure^[1].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p> |

REFERENCES

[1]. Kennedy GL Jr, et al. Inhalation toxicity of dehydrothio-p-toluidine. Food Chem Toxicol. 1984 Apr;22(4):289-92.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA