



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Sodium carboxymethyl cellulose (Viscosity:300-800mpa.s)

Cat. No.:	HY-Y0703C	
CAS No.:	9004-32-4	
Target:	Biochemical Assay Reagents	
Pathway:	Others	Sodium carboxymethyl cellulose
Storage:	4°C, sealed storage, away from moisture	
	* In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture)	

SOLVENT & SOLUBILITY

In Vitro	H ₂ O : 25 mg/mL (ultrasonic and warming and heat to 60°C)
----------	---

BIOLOGICAL ACTIVITY

Description	Sodium carboxymethyl cellulose (Viscosity:300-800mpa.s) is the sodium salt of cellulose arboxymethyl and frequently used as viscous agent, paste and barrier agent.
-------------	---

In Vivo	<p>Protocol for preparing 0.5% CMC-Na Solution Measure 0.5g of dry CMC-Na and dissolved in 100 ml ddH₂O/0.9% Saline (0.9 g NaCl in 100 ml ddH₂O) to make a clear solution. Under the condition of stirring and heating (50-65°C), adding CMC-Na slowly to ddH₂O/0.9% Saline helps to accelerate dissolution.</p> <p>Note</p> <ol style="list-style-type: none"> You must ensure that your CMC-Na solution does not exist solid-liquid separation phenomenon. The solution is in a uniform and transparent state has no particles in it. Completely dissolution of CMC-Na may requires 4 hours or more longer. <p>In a pharmacological test, CMC-Na (oral;5% in water; 1 year) is well tolerated in rats^[2]. In an acute oral toxicity study in female mice, LD₅₀ of CMC-Na for female mice is 14 g/kg body weight of mice, equivalent to 9.8 g/kg body weight of rat, categorized as practically non-toxic according to Loomis criteria (LD₅₀ 5-15g/kg body weight of rat)^[3]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>
---------	---

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA