



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## o-Vanillin

Cat. No.:	HY-Y1832
CAS No.:	148-53-8
Molecular Formula:	C <sub>8</sub> H <sub>8</sub> O <sub>3</sub>
Molecular Weight:	152.15
Target:	Fungal
Pathway:	Anti-infection
Storage:	4°C, stored under nitrogen
	* In solvent : -80°C, 6 months; -20°C, 1 month (stored under nitrogen)



### SOLVENT & SOLUBILITY

#### In Vitro

DMSO : 100 mg/mL (657.25 mM; Need ultrasonic)

Preparing Stock Solutions	Solvent Concentration	Mass		
		1 mg	5 mg	10 mg
	1 mM	6.5725 mL	32.8623 mL	65.7246 mL
	5 mM	1.3145 mL	6.5725 mL	13.1449 mL
	10 mM	0.6572 mL	3.2862 mL	6.5725 mL

Please refer to the solubility information to select the appropriate solvent.

### BIOLOGICAL ACTIVITY

#### Description

o-Vanillin (2-Vanillin) is a nature product, could be extracted from *Vanilla planifolia*, *Pinus koraiensis* fruit. o-Vanillin is a potent antifungal agent. o-Vanillin inhibits the growth of mycelia by disrupting the integrity of cell walls and cell membranes. o-Vanillin inhibits [Doxorubicin](#) (HY-15142A)- and 4-hydroperoxycyclophosphamide-induced NF-κB activation<sup>[1]</sup> [2].

#### In Vitro

o-Vanillin (2-Vanillin; 0-125 μg/mL; 24-72 h) inhibits the mycelial growth of *A. flavus* in a dose-dependent manner<sup>[1]</sup>.  
 o-Vanillin (0-100 μg/mL; 48 h; *A. flavus*) changes the morphology of mycelia and induces irregular shrinkage of the mycelia<sup>[1]</sup>.  
 o-Vanillin (0-100 μg/mL; *A. flavus*) decreases the protein content of the cell wall surface and the content of β-1,3-glucan<sup>[1]</sup>.  
 o-Vanillin (0-100 μg/mL; *A. flavus*) destroys cell membrane integrity. o-Vanillin releases cell constituents and decreases extracellular pH value<sup>[1]</sup>.  
 o-Vanillin (0-100 μg/mL) could effectively inhibit the growth of *A. flavus* on corn kernels<sup>[1]</sup>.  
 o-Vanillin (0-250 μM) inhibits doxorubicin-mediated induction of NFκB activity by 65% in A375/NFκB-Luc cells. o-Vanillin suppresses 4-HC-induced activity by 43%<sup>[2]</sup>.  
 MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### In Vivo

o-Vanillin (2-Vanillin; 60 mg/kg; p.o.; daily, for 5 d) inhibits tumor growth in mice bearing A375 human melanoma xenografts [2].

---

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	Male NSG mice with A375 human melanoma xenografts (12-16 weeks of age) <sup>[2]</sup>
Dosage:	60 mg/kg
Administration:	Oral administration; daily, for 5 days
Result:	Delayed the growth of A375 human melanoma xenografts in immunodeficient NSG mice.

---

## REFERENCES

[1]. Li Q, et, al. o-Vanillin, a promising antifungal agent, inhibits *Aspergillus flavus* by disrupting the integrity of cell walls and cell membranes. *Appl Microbiol Biotechnol*. 2021 Jun;105(12):5147-5158.

[2]. Marton A, et, al. Vanillin Analogues o-Vanillin and 2,4,6-Trihydroxybenzaldehyde Inhibit NF $\kappa$ B Activation and Suppress Growth of A375 Human Melanoma. *Anticancer Res*. 2016 Nov;36(11):5743-5750.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA