



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## (S)-2-[(tert-Butoxycarbonyl)amino]-3-iodopropionic acid methyl ester

<b>Cat. No.:</b>	HY-Z0424
<b>CAS No.:</b>	170848-34-7
<b>Molecular Formula:</b>	C <sub>9</sub> H <sub>16</sub> INO <sub>4</sub>
<b>Molecular Weight:</b>	329.13
<b>Target:</b>	Amino Acid Derivatives
<b>Pathway:</b>	Others
<b>Storage:</b>	4°C, protect from light * In solvent : -80°C, 6 months; -20°C, 1 month (protect from light)



### SOLVENT & SOLUBILITY

<b>In Vitro</b>	DMSO : 100 mg/mL (303.83 mM; ultrasonic and warming and heat to 60°C)				
		<b>Solvent</b>	<b>Mass</b>		
		<b>Concentration</b>	<b>1 mg</b>	<b>5 mg</b>	<b>10 mg</b>
	<b>Preparing Stock Solutions</b>	<b>1 mM</b>	3.0383 mL	15.1916 mL	30.3831 mL
		<b>5 mM</b>	0.6077 mL	3.0383 mL	6.0766 mL
<b>10 mM</b>		0.3038 mL	1.5192 mL	3.0383 mL	
Please refer to the solubility information to select the appropriate solvent.					
<b>In Vivo</b>	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (7.60 mM); Clear solution  2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (7.60 mM); Clear solution				

### BIOLOGICAL ACTIVITY

<b>Description</b>	(S)-2-[(tert-Butoxycarbonyl)amino]-3-iodopropionic acid methyl ester is an alanine derivative <sup>[1]</sup> .
<b>In Vitro</b>	Amino acids and amino acid derivatives have been commercially used as ergogenic supplements. They influence the secretion of anabolic hormones, supply of fuel during exercise, mental performance during stress related tasks and prevent exercise induced muscle damage. They are recognized to be beneficial as ergogenic dietary substances <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### REFERENCES

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA