



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Mouse anti double-stranded RNA (K2)

 nordicmubio.com/products/mouse-anti-double-stranded-rna-k2/10030005

Catalog number: **10030005**

Clone	K2
Isotype	IgM kappa
Product Type	Monoclonal Antibody
Units	5 ml
Host	Mouse
Application	ELISA Immunofluorescence

Background

Over the past decade our double-stranded RNA (dsRNA) antibodies have been used extensively to detect and characterise plant and animal viruses with dsRNA genomes or intermediates. In addition, the anti-dsRNA antibodies can be used as a diagnostic tool to detect pathogens, including detection in paraffin-embedded fixed tissue samples (Richardson et al. 2010). K2 anti-dsRNA monoclonal antibody has an IgM isotype. K2 has primarily been used in ELISA and sandwich ELISA. Generally, K2 can be used in applications where an anti-dsRNA antibody with an isotype other than IgG (IgG2a) is required.

Synonyms: Mouse anti dsRNA

Source

Female DBA/2 mice were injected intraperitoneally with a mixture of 50 ug L-dsRNA and 75 ug methylated bovine serum albumin, emulsified in complete Freund's adjuvant. After several boosts spleen cells were fused with Sp2/0-Ag14 myeloma cells to generate the hybridoma clone.

Product

The mAb K2 recognises double-stranded RNA (dsRNA) provided that the length of the helix is greater than or equal to 40 bp dsRNA. Recognition is independent of the sequence and nucleotide composition of the antigen. All naturally occurring dsRNAs

investigated up to now (4050 species) as well as poly(I).poly(C) and poly(A).poly(U) have been recognised by K2. As described by Schönborn et al. K2 binds with high avidity to all dsRNAs investigated.

Formulation: Culture supernatant (RPMI, 5% fetal calf serum).

Concentration: Undiluted hybridoma supernatant

Applications

MAB K2 is primarily used for a sandwich ELISA to detect and quantitate (after calibration) dsRNA (see Schönborn et al.). For this application it should be diluted 1:2 with PBS. It may also be advantageous to use K2 for immunofluorescence studies. The optimum working dilution of the antibody for any specific application should be established by titration. Not for use for clinical purposes. For in vitro use only.

Storage

After delivery antibodies should be aliquoted and stored at -20 ° or -70 °C. After adding 10 mM sodium azide undiluted antibody can also be stored at +4 °C for a short period of time. For long term storage the antibody should be kept frozen. Repeated freezing/thawing cycles should be avoided.

Shipping Conditions: The IgM monoclonal antibody K2 is only sold as hybridoma supernatant and shipped at ambient temperature.

Caution

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. It may contain hazardous ingredients. Please refer to the Safety Data Sheets (SDS) for additional information and proper handling procedures. Dispose product remainders according to local regulations. This datasheet is as accurate as reasonably achievable, but Exalpha Biologicals accepts no liability for any inaccuracies or omissions in this information.

References

1) Schönborn, J., Oberstrass, J., Breyel, E., Tittgen, J., Schumacher, J. and Lukacs, N. (1991) Monoclonal antibodies to double-stranded RNA as probes of RNA structure in crude nucleic acid extracts. *Nucleic Acids Res.* 19, 2993-3000. 2) Lukacs, N. (1994) Detection of virus infection in plants and differentiation between coexisting viruses by monoclonal antibodies to double-stranded RNA. *J. Virol. Methods* 47, 255-272. 3) Lukacs, N. (1997) Detection of sense:antisense duplexes by structurespecific anti-RNA antibodies. In: *Antisense Technology. A Practical Approach*, C. Lichtenstein and W. Nellen (eds), pp. 281-295. IRL Press, Oxford. 4) S. J. Richardson, A. Willcox, D. A. Hilton, S. Tauriainen, H. Hyoty, A. J. Bone, A. K. Foulis, N. G. Morgan. Use of antisera directed against dsRNA to detect viral infections in formalin-fixed paraffin-embedded tissue. *J Clin Virol.* (2010) 49(3); 180-5. doi: 10.1016/j.jcv.2010.07.015.

