



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



## Goat anti Human IgG Fab, conjugated with Biotin

Catalogue number: **GAHu/Fab/Bio**

Clone	Polyclonal
Product Type	Secondary Antibodies
Units	1 ml
Host	Goat
Species reactivity	Human
Application	Dot blot ELISA Immunoblotting Immunocytochemistry Immunohistochemistry (paraffin)

### Background

The antiserum is reacting with the Fab subunit of intact IgG, IgA and IgM and other Ig classes of both light chain types, with their Fab or F(ab')<sub>2</sub> subunits and with free light chains of kappa and lambda type as tested in immunoelectrophoresis and double radial immunodiffusion. In immunocytochemical and immunohistochemical staining to identify and measure free or Ig bound subunits of both kappa and lambda type at the cellular and subcellular level by staining of appropriately treated cell and tissue substrates, and to demonstrate circulating antibodies in serodiagnostic microbiology and autoimmune diseases; to identify a specific antigen or immune complex using a reference antibody of human origin in the middle layer of the indirect test procedure. As a second step an avidin or streptavidin conjugate of the user's choice has to be used. In general this kind of products is not recommended as direct or indirect screening reagents for immunoglobulin isotypes on the surface of membranes of vital lymphoid cells. The presence of activity to the common Fab subunit may result in the staining of Ig bound to Fc-receptors on non-lymphoid cells. Combinations of isotype-specific reagents should be used instead for this purpose. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal.

### Source

Purified Fab from normal IgG isolated from pooled human serum.

## Distributors

For Purchasing Information,  
please contact your local  
distributor

[Find Distributor](#)

Freund's complete adjuvant is used in the first step of the immunization procedure.

**Product**

Biotin-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added.

**Applications**

ELISA, Immunocytochemistry, Immunohistochemistry (paraffin), Dot blot, Immunoblotting.

**Cross Reactivity**

Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this antiSerum has not been tested in detail.

**Specificity**

Biotin-conjugated IgG fraction of polyclonal Goat antiSerum to Human Fab of IgG

**Storage**

The lyophilized conjugate is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20°C. Prior to use, an aliquot is thawed slowly in the dark at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the immunoconjugate. Lyophilized at +4° C--at least 10 years. Reconstituted at or below -20° C--3-5 years. Reconstituted at +4° C--7 days

**Caution**

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Nordic-MUbio accepts no liability for any inaccuracies or omissions in this information.