



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Goat anti Human fibrinogen, conjugated with Biotin

Catalogue number: **GAHu/Fbg/Bio**

Clone	Polyclonal
Product Type	Primary Antibodies
Units	1 ml
Host	Goat
Species reactivity	Human
Application	Dot blot ELISA Immunoblotting Immunocytochemistry Immunohistochemistry (paraffin)

Background

The reactivity of the antiserum is restricted to fibrinogen. In immunoelectrophoresis and radial immunodiffusion (Ouchterlony), using various antiserum concentrations against normal human plasma a single precipitin line is obtained which shows a reaction of identity with the precipitin line obtained with purified fibrinogen. No reaction is obtained with any other plasma protein component or serum. However, the antiserum may also react with fibrin monomers, circulating fibrinopeptides and fibrin degradation products. As reagent for the direct detection of fibrinogen in human cells, tissues and body fluids in immunocytochemical and immunohistochemical assays; as detection reagent in non-isotopic methodology and solid phase immunochemistry (e.g. ELISA). As a second step an avidin or streptavidin conjugate of the user's choice has to be used. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal.

Source

Fibrinogen (clotting factor I) is a heat labile beta glycoprotein present in plasma. It is the precursor of fibrin, which is the key protein constituting the network of the blood clot. Thrombin converts fibrinogen to fibrin by limited proteolysis. Fibrin monomers polymerize to fibrin which is stabilized by cross-linking. Fibrinogen is isolated from fresh plasma after removing prothrombin. Freund's complete adjuvant is used in the first step

Distributors

For Purchasing Information,
please contact your local
distributor

[Find Distributor](#)

of the immunization procedure.

Product

Biotin-coupled purified hyperimmune IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2) No preservative added, as it may interfere with the antibody activity.

Applications

ELISA, Immunocytochemistry, Immunohistochemistry (paraffin), Dot blot, Immunoblotting. The lyophilized conjugate is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20°C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided.

Cross Reactivity

The antiSerum does not cross-react with any other component of Human plasma. Inter-species crossreactivity is a normal feature of antibodies to plasma proteins since they frequently share antigenic determinants. Cross-reactivity of this antiSerum with Bovine, Dog, Cat and Monkey has been observed.

Specificity

Biotin-conjugated IgG fraction of polyclonal Goat antiSerum to Human fibrinogen

Storage

Lyophilized at +4°C--at least 10 years. econstituted at or below -20°C--3-5 years. econstituted at +4°C--7 days

Caution

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Nordic-MUbio accepts no liability for any inaccuracies or omissions in this information.