



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Goat anti Human IgG (Fc specific) immunofixation

Catalogue number: **GAHu/IgG(Fc)/IFix**

Clone	Polyclonal
Product Type	Secondary Antibodies
Units	1 ml
Host	Goat
Species reactivity	Human
Application	Immunoprecipitation

Background

The reactivity of the antiserum is directed to the Fc subunit of the IgG molecule which expresses strict isotypic (class) specificity. It does not react with any non-Ig protein in human serum, as tested by immunoelectrophoresis and double radial immunodiffusion. This antiserum is intended to detect and identify IgG in serum or other body fluids using the immunofixation technique. Like immunoelectrophoresis immunofixation is essentially a two step technique. Proteins in a complex mixture are separated by electrophoresis in a gel carrier, followed by immunoprecipitation in situ with the antiserum. Non-precipitated proteins are removed by washing and the precipitated complex is revealed with a protein stain which allows its exact localization. Immunofixation may be the method of choice whenever a high level of sensitivity is required to identify a minor protein component against a high background of other proteins. It enables the detection and identification of more than one paraproteins in serum or of free light chain. The detection limit is approximately 0.5 to 1 mg/ml in the presence of normal levels of immunoglobulins. This product is not pre-diluted. The optimum working dilution of each product should be established by titration before being used.

Source

Highly purified polyclonal IgG isolated from human serum. Freund's complete adjuvant is used in the first step of the immunization procedure.

Product

Goat Ig lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity.

Applications

Distributors

For Purchasing Information,
please contact your local
distributor

[Find Distributor](#)

Immunoprecipitation.

Cross Reactivity

Inter-species cross-reactivity is a normal feature of antibodies to mammalian immunoglobulins, since homologous proteins of different species frequently share antigenic determinants. The degree of crossreactivity is also dependent on the concentrations of the reactants and the sensitivity of the assay arrangement. This antiSerum fraction has not been tested for cross-reactivity.

Specificity

Ig fraction of polyclonal Goat antiSerum to Human immunoglobulin IgG, Fc specific

Storage

The lyophilized product is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20°C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the product. Lyophilized at +4° C--at least 10 years. Reconstituted at or below -20° C--3-5 years. Reconstituted at +4° C--7 days

Caution

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Nordic-MUbio accepts no liability for any inaccuracies or omissions in this information.