



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Goat anti Human kininogen

Catalogue number: **GAHu/Kg**

Clone	Polyclonal
Product Type	Primary Antibodies
Units	1 ml
Host	Goat
Species reactivity	Human
Application	Immunoprecipitation

Distributors

For Purchasing Information, please contact your local distributor

[Find Distributor](#)

Background

The heavy chains in HMWK and LMWK share immunologic determinants, but they have no procoagulant activities. The light chain of HMWK is unique to this molecule and carries the entire procoagulant activity of the molecule. The antiserum does not distinguish between different forms of kininogen. The presence of non-precipitating antibodies has not been assayed. If the antiserum is to be used in non-precipitating antibody-binding assays (immunocytochemical, immunohistochemical, solid phase procedures such as ELISA) proper controls should be included. No reaction is obtained with any other plasma protein component or serum. In precipitating techniques as immunoelectrophoresis and single or double radial immunodiffusion and electroimmunodiffusion (Laurell), to identify the presence of kininogen in human plasma or other body fluids or to determine its concentration. For electroimmunodiffusion a concentration of 1-2% of antiserum in the gel is recommended.

Source

Human plasma contains at least two distinct forms of kininogen, the carrier molecule of bradykinin, which can be distinguished by their physical, chemical and immunochemical properties. They are designated high-molecular-weight kininogen (HMWK) and low-molecular weight kininogen (LMWK). HMWK (MW 110,000) and LMWK (MW 50,000) both liberate bradykinin following the action of kallikrein. HMWK circulates as a non-enzymatic factor that is central to contact activation reactions. LMWK plays no known role in the activation of coagulation. Normal adult plasma levels are 7 mg/100 ml for HMWK as determined by electroimmunodiffusion and 9 mg/100 ml based on haemagglutination inhibition assays and RIA. Low plasma kininogen levels are seen in newborn and pre-term infants. HMWK is absent in congenital kininogen deficiency of the named affected families Fitzgerald and Reid; Williams and Flaujeac lack both HMWK and LMWK. The condition is associated with the decreased fibrinolytic activity and kinin formation but is

asymptomatic. Purified kininogen from pooled human plasma is used for the immunization. Freund's complete adjuvant is used in the first step of the immunization procedure.

Product

Delipidated, heat inactivated, lyophilized, stable whole antiserum
No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum.
No foreign proteins added.

Applications

Immunoprecipitation. For electroimmunodiffusion a concentration of 1-2% of antiserum in the gel is recommended.

Cross Reactivity

The antiSerum does not cross react with any other component of Human plasma. Inter-species crossreactivity is a normal feature of antibodies to plasma proteins since they frequently share antigenic determinants. Cross-reactivity of this antiSerum has not been tested in detail.

Specificity

Precipitating polyclonal Goat antiSerum to Human kininogen

Storage

The lyophilized antiserum is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C.
Reconstitute the lyophilized antiserum by adding 1 ml sterile distilled water. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the antiserum. Diluted antiserum should be stored at +4°C, not refrozen, and preferably used the same day.
Lyophilized at +4°C--at least 10 years. Reconstituted at or below -20°C--3-5 years. Reconstituted at +4°C--7 days

Caution

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Nordic-MUBio accepts no liability for any inaccuracies or omissions in this information.