



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Goat anti Monkey IgG IgA IgM (Fc specific), conjugated with Biotin

Catalogue number: **GAMon/Ig(Fc)/Bio**

Clone	Polyclonal
Product Type	Secondary Antibodies
Units	1 ml
Host	Goat
Species reactivity	Monkey
Application	Dot blot ELISA Immunoblotting Immunocytochemistry Immunohistochemistry (paraffin)

Background

The reactivity of the antiserum is directed to the Fc subunit of the major immunoglobulin isotypes which expresses strict isotypic (class) specificity. In immunoelectrophoresis and double radial immunodiffusion using various antiserum concentrations against normal monkey plasma and serum, the characteristic IgG, IgA and IgM precipitin lines are obtained. In immunocytochemical and immunohistochemical staining of immunoglobulins at the cellular and subcellular level of appropriately treated cell and tissue substrates; to demonstrate circulating antibodies in serodiagnostic microbiology and autoimmune diseases; to identify a specific antigen using a reference antibody of monkey origin in the middle layer of the indirect test procedure; in non-isotopic assay methodology (e.g. ELISA) to measure immunoglobulins in monkey serum or other body fluids. The absence of activity to the common Ig/Fab subunit prevents the reaction of this conjugate with immunoglobulins binds to Fc receptors on non-lymphoid cells. As a second step an avidin or streptavidin conjugate of the user's choice has to be used. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal.

Source

Purified normal IgG, IgA and IgM isolated from pooled rhesus monkey serum. Freund's complete adjuvant is used in the first step of the immunization procedure.

Distributors

For Purchasing Information, please contact your local distributor

[Find Distributor](#)

Product

Biotin-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity.

Applications

ELISA, Immunocytochemistry, Immunohistochemistry (paraffin), Dot blot, Immunoblotting.

Cross Reactivity

Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Precipitation reactions have been observed with immunoglobulins in Serum of other old-world Monkeys, including Cercopithecus, Cynomolgus and Baboon. The conjugate may also react with other species as has been observed for Chimpanzee and man.

Specificity

Biotin-conjugated IgG fraction of polyclonal Goat antiSerum to Monkey IgG, IgA and IgM, Fc specific.

Storage

The lyophilized conjugate is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20°C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the immunoconjugate. Lyophilized at +4° C--at least 10 years. Reconstituted at or below -20° C--3-5 years. Reconstituted at +4° C--7 days

Caution

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Nordic-MUbio accepts no liability for any inaccuracies or omissions in this information.