



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



## Goat anti Monkey IgG IgA IgM (Fc specific), conjugated with FITC

Catalogue number: **GAMon/Ig(Fc)/FITC**

Clone	Polyclonal
Product Type	Secondary Antibodies
Units	1 ml
Host	Goat
Species reactivity	Monkey
Application	ELISA Immunocytochemistry Immunofluorescence Immunohistochemistry (frozen)

## Distributors

For Purchasing Information, please contact your local distributor

[Find Distributor](#)

### Background

The reactivity of the antiserum is directed to the Fc subunits of the major isotypes of the monkey immunoglobulin system. No reaction is obtained with purified IgG/Fab or any non-Ig protein of monkey serum, as tested by immunoelectrophoresis and double radial immunodiffusion. In immunoelectrophoresis and double radial immunodiffusion using various antiserum concentrations against normal monkey plasma and serum, the characteristic IgG, IgA and IgM precipitin lines are obtained. Direct immunofluorescence staining of cytoplasmic Ig of appropriately treated cell and tissue substrates; to demonstrate immunoglobulins or specific antibodies in cells and tissues; to identify circulating antibodies in serodiagnostic microbiology and autoimmune diseases; to identify a specific antigen or immune complex using a reference antibody of monkey origin in the middle layer of the indirect test procedure. The absence of activity to the common Ig/Fab subunit prevents the reaction of this conjugate with immunoglobulins binds to Fc receptors on non-lymphoid cells. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal.

### Source

Purified polyclonal monkey IgG, and IgA and IgM containing fractions isolated from monkey serum. Freund's complete adjuvant is used in the first step of the immunization procedure.

### Product

FITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity.

#### **Applications**

ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence.

#### **Cross Reactivity**

Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Precipitation reactions have been observed with immunoglobulins in Serum of other old-world Monkeys, including Cercopithecus, Cynomolgus and Baboon. The conjugate may also react with other species as has been observed for Chimpanzee and man.

#### **Specificity**

Fluorescein isothiocyanate-conjugated IgG fraction of polyclonal Goat antiSerum to Monkey IgG, IgA and IgM, Fc specific.

#### **Storage**

The lyophilized conjugate is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. It is reconstituted by adding 1 ml sterile distilled water. Spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20°C. Prior to use, an aliquot is thawed slowly in the dark at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the immunoconjugate. Lyophilized at +4° C--at least 10 years. Reconstituted at or below -20° C--3-5 years. Reconstituted at +4° C--7 days

#### **Caution**

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Nordic-MUbio accepts no liability for any inaccuracies or omissions in this information.